arXiv:1110.1711v2 [math.CT] 24 Aug 2012

Inner products of 2-representations

Nora Ganter*
The University of Melbourne

August 27, 2012

Abstract

We define and calculate inner products of 2-representations. Along the way, we prove that the categorical trace $\mathbb{T}r(-)$ of [GK08, Sec.3] is multiplicative with respect to various notions of categorical tensor product, and we identify the center of the category of equivariant objects $\mathcal{V}^G$ of [GK08, Sec.4.2]. We discuss applications, ranging from Schur's result about the number of irreducible projective representations to a formula for the Hochschild cohomology of a global quotient orbifold.

Contents

1 Introduction and statement of results **2**
1.1 String diagram notation 6

2 Multiplicativity of the categorical trace **8**
2.1 The multiplicativity theorem 9
2.2 Examples 13
2.2.1 2-Vector spaces 13
2.2.2 Algebras 13
2.2.3 Coherent sheaves 15

---

[*] Ganter was supported by NSF-grant DMS-0504539 and by a Centenary Fellowship by the Faculty of Science at the University of Melbourne. Work on this project began while Ganter was at Colby College, Maine.

3 Traces in monoidal 2-categories **16**
3.1 Bimodules, strict version 16
3.2 Bimodules, derived version 17
3.3 Constructible sheaves 17
3.4 Coherent sheaves 19

4 Tensor products of 2-representations **23**

5 Categories of equivariant objects **26**
5.1 Background and Definitions 26
5.2 The twisted group algebra 32
5.3 The G-action on the twisted group algebra 35
5.4 Traces in V^G 37
5.5 Inner products 39

6 Applications **39**
6.1 Projective representations 39
6.2 Algebras 40
6.3 Bimodules 41
6.4 Coherent sheaves 42
6.5 The example of a trivial G-action 44

A Clean intersections **45**

1 Introduction and statement of results

The idea of a *representation of a group by functors in a category* goes back to Grothendieck [Gro57, p.196], and categorified versions of representation theory have since been developed by a number of authors.[1] (We recall some of the basic definitions in Section 4.) The corresponding character theory was introduced in [Bar08] and [GK08]. Key ingredient of this categorified character theory is the notion of categorical trace: let F be an endofunctor of a small category. Then the *categorical trace* of F is defined as the set of natural transformations

$$\mathbb{T}r(F) = \mathcal{N}\mathrm{at}(\mathrm{id}, F).$$

---

[1] See for instance [Elg07], [CY05], [BM06], [Ost03], [Fre94], [BWC07], [Del97], [ST01], [BZ07] or [Kho00].

More generally, if F is a 1-endomorphism in a 2-category, then

$$\mathbb{T}r(F) = 2\operatorname{Hom}(1, F).$$

Let ϱ be a linear 2-representation of a finite group G. Then its *categorical character* X_ϱ sends $g \in G$ to the k-vector space

$$X_\varrho(g) = \mathbb{T}r(\varrho(g)).$$

The categorical character of ϱ comes equipped with conjugation isomorphisms

$$\psi_h \colon X_\varrho(g) \xrightarrow{\cong} X_\varrho(h^{-1}gh),$$

one for each pair of elements of G (see [GK08, Prop.4.10]). As g and h vary, the ψ_h are compatible, endowing X_ϱ with the structure of a *categorical class function* on G. Such categorical class functions were first introduced by Luztig [Lus87], who named them *class sheaves*.

For a commuting pair (g, h), the map ψ_h is an automorphism of $X_\varrho(g)$. Assume that all the $X_\varrho(g)$ are finite dimensional. Then the *2-character* χ_ϱ of ϱ is the function

$$\chi_\varrho(g, h) = tr(\psi_h),$$

defined on pairs of commuting elements of G. It satisfies

$$\chi_\varrho(s^{-1}gs, s^{-1}hs) = \chi_\varrho(g, h).$$

Such conjugation invariant functions on commuting pairs are called *2-class functions*.

Much of our motivation for writing [GK08] came from a character theory discovered by Hopkins, Kuhn and Ravenel in the context of stable homotopy theory [HKR00]. Although of entirely different origin, these Hopkins-Kuhn-Ravenel characters exhibit strikingly similar features to ours. For instance, we proved in [GK08, Cor.6.12] that the 2-character of an induced 2-representation is described by the same formula as the effect of transfer on Hopkins-Kuhn-Ravenel characters.

In this paper we will define inner products of 2-representations and calculate them in terms of categorical characters and 2-characters. On the level of 2-characters we obtain

$$\langle \chi, \xi \rangle_G = \frac{1}{|G|} \sum_{gh=hg} \chi(g, h) \cdot \xi(g, h) \tag{1}$$

(see Corollary 5.18). This agrees with the formula for the Strickland inner product of Hopkins-Kuhn-Ravenel characters in [Gan06, Prop.1.6]. Our definition will be the categorical analogue of the inner product

$$\langle V, W\rangle_G := \dim_k (V \otimes W)^G$$

on the representation ring $R(G)$ (compare [Str00]). This is closely related to the usual inner product, sending (V, W) to

$$\dim_k (\mathrm{Hom}_G(W, V)) = \langle V, W^*\rangle_G.$$

The paper is organized as follows: in Section 2, we let g and h be endofunctors of linear categories $\mathcal{V}$ and $\mathcal{W}$, respectively. Under a finiteness condition we construct an isomorphism

$$\mu\colon \mathbb{T}r(g) \otimes \mathbb{T}r(h) \cong \mathbb{T}r(g \boxtimes h)$$

(Theorem 2.5). Here $\boxtimes$ is the tensor product of linear categories defined in [GK, Sec.1]. Similar isomorphisms are found in the abelian case (using the Deligne tensor product, [Del90]) and in the pre-triangulated case (using the Bondal-Larsen-Lunts tensor product [BLL04]). We then discuss examples. The map μ exists in the more general context of monoidal 2-categories. For some of those, μ is an isomorphism.

In Section 5 we let ϱ be an action of G on a linear category $\mathcal{V}$, and consider the category $\mathcal{V}^G$ of equivariant objects of ϱ. We introduce the *twisted group algebra*

$$R_\varrho = \bigoplus_{g\in G} \mathbb{T}r(g)$$

acting on each object of $\mathcal{V}^G$. We contruct an isomorphism of k-algebras

$$\mathcal{D}\mathrm{im}\left(\mathcal{V}^G\right) \cong R_\varrho^G. \tag{2}$$

Here

$$\mathcal{D}\mathrm{im} = \mathbb{T}r(1)$$

is the center (of the category $\mathcal{V}^G$). As a corollary, we obtain

$$\dim\left(\mathcal{D}\mathrm{im}\left(\mathcal{V}^G\right)\right) = \frac{1}{|G|}\sum_{gh=hg} \chi_\varrho(g, h).$$

The category $\mathcal{V}^G$ is characterized by a universal property, which makes sense in the 2-categorical setup. All our proofs go through in the context of 2-categories.

Given two linear 2-representations V and W of G, we can now define their inner product as

$$\langle V, W\rangle_G = \mathcal{D}\mathrm{im}\,(V \boxtimes W)^G\,,$$

and conclude that, if $\langle V, W\rangle_G$ exists, its k-dimension is calculated by (1). In Section 6, we discuss applications of (2). One special case is a Theorem by Schur, counting the isomorphism classes of irreducible projective representations with a given cocycle c. Another special case is as follows: let X be a smooth projective variety over $\mathbb{C}$, acted upon by G. Let ϱ be the resulting 2-representation of G in $\mathcal{V}\mathrm{ar}_{\mathbb{C}}$ (see [GK08, 2.4.(c)]). In this context (2) becomes a result about orbifold Hochschild cohomology, namely

$$\mathrm{HH}^\bullet([X/G]) \cong \bigoplus_{[g],\alpha} \mathrm{HH}^{\bullet\text{-}\mathrm{codim}(\mathrm{X}^{\mathrm{g}}_\alpha)}\,(X^g_\alpha, \det N^g)\,. \tag{3}$$

Here X^g_α are the connected components of the fixed point set X^g of g, and N^g is the normal bundle of X^g in X; further,

$$\mathrm{HH}^\bullet(Y, \mathcal{F}) \coloneqq \mathrm{Ext}_{\mathrm{Y}\times\mathrm{Y}}(\Delta_*\mathscr{O}_Y, \Delta_*\mathcal{F}).$$

I understand that an additive version of the isomorphism (3), with a very different proof, was known to Andrei Caldararu.

The main advantage of our approach is multiplicativity: the right-hand side of (3) is canoically an algebra, namely the twisted group algegra of the 2-representation ϱ, and (3) is an algebra map. Similar isomorphisms can be found in [Căl05], [DE05], [GK04] and [Bar03], and it is a common theme that the proof of multiplicativity tends to be quite difficult. We note that our proof of that fact is very simple (see page 38).

Some interesting questions about multiplicativity remain, for instance, whether it is possible to give an explicit formula for the product on the right-hand side, involving the double fixed point sets $X^g \cap X^h$.

Further applications of our results turn up in [GK], where we define symmetric and exterior powers of categories and calculate their effects on characters of 2-representations.

Acknowledgements

This paper is my account of results that were, to a large extent, found in collaboration with Mikhail Kapranov. It is part of a series of sequels to [GK08]. Many crucial ideas were Mikhail's, and his comments on older drafts have been very helpful. It is a pleasure to thank him for all this. I would also like to thank Alex Ghitza, Ezra Getzler, and Craig Westerland for numerous helpful conversations on the topic.

1.1 String diagram notation

We will use string diagram notation for 2-morphisms. For a nice, short introduction to string diagram notation, we refer the reader to [CW10], for a more comprehensive account, we refer the reader to [Bar08]. What follows is a mini summary of what we will use. Our conventions follow those of Caldararu and Willerton. Bartlett's conventions are slightly different.[2]



Traditional versus string diagram notation for a 2-morphism $\phi\colon f \Rightarrow g$ between 1-morphisms $f, g\colon X \to Y$.

String diagrams are Poincare dual to the classical notation. They are read from the right to the left (direction of 1-morphisms) and from the bottom to the top (direction of 2-morphisms). The objects X and Y are sometimes omitted from the picture, when they are clear from the context.

Identity morphisms are denoted by omission:

---

[2] In his diagrams 2-morphisms flow downwards, while ours flow upwards.



A 2-morphism $\phi\colon \mathrm{id}_X \Rightarrow f$



A 2-morphism $\phi\colon f \Rightarrow \mathrm{id}_X$



The identity 2-morphism id_f

Composition is denoted by juxtaposition:



Vertical composition $\psi \circ_1 \phi$



Two ways of drawing the horizontal composition of 2-morphisms, $\psi\phi\colon hf \Rightarrow kg$. The fact that both diagrams describe the same 2-morphisms is known as the "butterfly identity" $(\psi g) \circ_1 (h\phi) = (k\phi) \circ_1 (\psi f) =: \psi\phi$.

Definition 1.1. An *adjunction* in a 2-category $\mathcal{C}$ consists of the following data: two objects, X and Y, 1-morphisms $f\colon X \to Y$ and $g\colon Y \to X$ and 2-morphisms



$\eta\colon \mathrm{id}_X \Rightarrow gf$, called the *unit* and denoted by either of these two pictures



and $\epsilon\colon fg \Rightarrow \mathrm{id}_Y$, called the *counit* and denoted by either of these two pictures.

These are required to satisfy



$(\mathrm{id}_g\,\epsilon)(\eta\,\mathrm{id}_g) = \mathrm{id}_g$, in other words, the 2-morphisms in these two string diagrams are required to be equal,



and $(\epsilon\,\mathrm{id}_f)(\mathrm{id}_f\,\eta) = \mathrm{id}_f$, i.e., the 2-morphisms in these two string diagrams are required to be equal.

We say "f is left adjoint to g".

For more on adjunctions in 2-categories, see for instance [CW10, p. 13 ff.].

2 Multiplicativity of the categorical trace

Let k be an algebraically closed field. By a *linear category*, we will always mean an additive k-linear category, and by a *linear functor* we mean one that preserves this structure. In [GK], we consider three different notions of tensor product of categories:

1. if $\mathcal{V}$ and $\mathcal{W}$ are linear categories, we consider the uncompleted tensor product $\mathcal{V} \boxtimes \mathcal{W}$, following [KV94] and [BK01],

2. if $\mathcal{A}$ and $\mathcal{B}$ are abelian linear categories, we consider the completed version $\mathcal{A}\widehat{\boxtimes}\mathcal{B}$, following Deligne [Del90],
3. if $\mathcal{C}$ and $\mathcal{D}$ are perfect dg-categories over k, we consider the completed tensor product $\mathcal{C}\widehat{\boxtimes}\mathcal{D}$, following [BLL04].

The goal of this section is to find isomorphisms

$$\mu\colon \mathbb{T}r(g) \otimes \mathbb{T}r(h) \cong \mathbb{T}r(g \boxtimes h) \cong \mathbb{T}r(g \widehat{\boxtimes} h),$$

where g and h are endofunctors of the relevant categories. We can always define a map μ, but in order to show that it is an isomorphism, we need a finiteness condition.

2.1 The multiplicativity theorem

The first two constructions above are characterized by similar universal properties. For instance, the data of (1) consist of a linear category $\mathcal{V}\boxtimes\mathcal{W}$ together with a k-bilinear functor

$$\boxtimes\colon \mathcal{V} \times \mathcal{W} \longrightarrow \mathcal{V} \boxtimes \mathcal{W},$$

which is universal in the following sense: let $\mathcal{Z}$ be a third linear category. Then precomposition with $\boxtimes$ defines an equivalence of categories

$$\mathcal{F}\mathrm{un}_{\mathrm{lin}}(\mathcal{V} \boxtimes \mathcal{W}, \mathcal{Z}) \xrightarrow{\simeq} \mathcal{F}\mathrm{un}_{\mathrm{bil}}(\mathcal{V} \times \mathcal{W}, \mathcal{Z}).$$

Here $\mathcal{F}\mathrm{un}_{\mathrm{lin}}$ and $\mathcal{F}\mathrm{un}_{\mathrm{bil}}$ stand for the categories of k-linear and k-bilinear functors.

The categorical tensor products in (1) and (3) always exist. In the abelian case, (2), Deligne has proved the existence of $\mathcal{A}\widehat{\boxtimes}\mathcal{B}$ under the following finiteness condition:

Definition 2.1 (compare [Del90, (2.12.1)])**.** We say that a linear abelian category $\mathcal{A}$ *satisfies Deligne's finiteness condition*, if all objects of $\mathcal{A}$ have finite length and all Hom-sets are finite dimensional k-vector spaces.

Kapranov and Voevodsky argued that $\boxtimes$ endows the 2-category $\mathcal{C}\mathrm{at}_{\mathrm{k}}$ of k-linear categories with the structure of a monoidal 2-category.The reader interested in the full twelve pages of axioms is referred to [KV94].[3] The following proposition, which follows immediately from the universal propetry, summarizes what we will need.

---

[3] The definition was later revised by Crans [Cra98], see also [BL03].

Proposition 2.2. *(1) Let $\mathcal{V}$, $\mathcal{V}'$, $\mathcal{W}$ and $\mathcal{W}'$ be linear categories. Then there exist a bilinear functor*

$$(- \boxtimes -)\colon \mathcal{F}\mathrm{un}_{\mathrm{lin}}(\mathcal{V},\mathcal{V}') \times \mathcal{F}\mathrm{un}_{\mathrm{lin}}(\mathcal{W},\mathcal{W}') \quad \longrightarrow \quad \mathcal{F}\mathrm{un}_{\mathrm{lin}}(\mathcal{V}\boxtimes\mathcal{W},\mathcal{V}'\boxtimes\mathcal{W}')$$

and a natural isomorphism

$$\iota\colon \boxtimes \circ (-,-) \Longrightarrow (- \boxtimes -) \circ \boxtimes.$$

The pair $((- \boxtimes -),\iota)$ is unique up to unique natural isomorphism (preserving ι). We have canonical functor isomorphisms

$$\kappa_{g,f,h,k}\colon (gf) \boxtimes (hk) \xrightarrow{\cong} (g \boxtimes h)(f \boxtimes k),$$

whenever the compositions on the left-hand side exist. They are natural in g, f, h, and k. Further, there is a functor isomorphism

$$\kappa_1\colon \mathrm{id}_{\mathcal{V}} \boxtimes \mathrm{id}_{\mathcal{W}} \xrightarrow{\cong} \mathrm{id}_{\mathcal{V}\boxtimes\mathcal{W}}.$$

These $\kappa_?$ are compatible with ι in an obvious sense and make the usual associativity pentagon and unit diagrams commute.

(2) Let $(\mathcal{V}\widetilde{\boxtimes}\mathcal{W},\widetilde{\boxtimes})$ be a second pair satisfying the universal property of $(\mathcal{V}\boxtimes\mathcal{W},\boxtimes)$. Then there exist an equivalence

$$E\colon \mathcal{V}\boxtimes\mathcal{W} \longrightarrow \mathcal{V}\widetilde{\boxtimes}\mathcal{W}$$

and a natural isomorphism

$$\iota_E\colon \widetilde{\boxtimes} \Rightarrow E \circ \boxtimes.$$

The pair (E,ι_E), is unique up to unique natural isomorphism(preserving ι_E).

A similar statement holds for the abelian case and right-exact linear functors.

Definition 2.3. Let $\mathcal{V}$ and $\mathcal{W}$ be linear categories, let g be an endofunctor of $\mathcal{V}$, and let h be an endofunctor of $\mathcal{W}$. Then the proposition gives a bilinear map

$$\widetilde{\mu}\colon \mathbb{T}r(g) \times \mathbb{T}r(h) \longrightarrow \mathcal{N}\mathrm{at}(\mathrm{id}_{\mathcal{V}} \boxtimes \mathrm{id}_{\mathcal{W}}, g \boxtimes h) \longrightarrow \mathbb{T}r(g \boxtimes h)$$
$$(\phi,\psi) \longmapsto (\phi \boxtimes \psi) \circ \kappa_1^{-1}.$$

We let μ be the resulting linear map

$$\mu\colon \mathbb{T}r(g) \otimes \mathbb{T}r(h) \longrightarrow \mathbb{T}r(g \boxtimes h).$$

Note that μ is natural in g and h. The following is a reformulation of [Del90, Prop.5.13 (i),(v)]:

Proposition 2.4. *Let $\mathcal{A}$ and $\mathcal{B}$ be k-linear abelian categories satisfying Deligne's finiteness condition. Then the completed tensor product $\mathcal{A}\widehat{\boxtimes}\mathcal{B}$ exists. Further, there is a fully faithful "completion" functor*

$$\mathcal{A} \boxtimes \mathcal{B} \longrightarrow \mathcal{A}\widehat{\boxtimes}\mathcal{B},$$

obtained by applying the universal property of $\boxtimes$ to $\widehat{\boxtimes}$.

In particular, the essential image of $\widehat{\boxtimes}\colon \mathcal{A} \times \mathcal{B} \to \mathcal{A}\widehat{\boxtimes}\mathcal{B}$ is contained in $\mathcal{A}\boxtimes\mathcal{B}$, and the two functors $\widehat{\boxtimes}$ and $\boxtimes$ are identified when viewed as functors onto their essential images.

By construction, there is a similar fully faithful "completion functor" for the Bondal-Larsen-Lunts tensor product of pre-triangulated categories.

Theorem 2.5. *(1) Let $\mathcal{V}$ and $\mathcal{W}$ be linear categories such that all* Hom*-sets are finite dimensional k-vector spaces. Let g be a linear endofunctor of $\mathcal{V}$, and let h be a linear endofunctor of $\mathcal{W}$. Assume that either $\mathbb{T}r(g)$ or $\mathbb{T}r(h)$ is finite dimensional. Then the map μ of Definition 2.3 is an isomorphism.*

(2) Let $\mathcal{A}$ and $\mathcal{B}$ be linear abelian categories satisfying Deligne's finiteness condition. Let g be a right-exact endofunctor of $\mathcal{A}$, and let h be a right-exact endofunctor of $\mathcal{B}$. Assume that either $\mathbb{T}r(g)$ or $\mathbb{T}r(h)$ is finite dimensional. Then we have an isomorphism

$$\mathbb{T}r(g \boxtimes h) \longrightarrow \mathbb{T}r(g\widehat{\boxtimes}h),$$

which is natural in g and h.

Proof : Part (2): let $\mathcal{A}$, $\mathcal{B}$, g and h be as in (2). Then we have

$$\begin{aligned} \mathbb{T}r(g\widehat{\boxtimes}h) &\cong \mathcal{N}\mathrm{at}(\widehat{\boxtimes}, \widehat{\boxtimes} \circ (g \times h)) \\ &= \mathcal{N}\mathrm{at}(\boxtimes, \boxtimes \circ (g \times h)) \\ &\cong \mathbb{T}r(g \boxtimes h). \end{aligned}$$

Here the first and third equality come from the universal properties of $\widehat{\boxtimes}$ and $\boxtimes$. The key step is the second equality, which follows from Proposition 2.4.

Part (1): Let (X, Y) be an object of $\mathcal{V} \times \mathcal{W}$. By construction of $\mathcal{V} \boxtimes \mathcal{W}$, we have

$$\mathrm{Hom}_{\mathcal{V}\boxtimes\mathcal{W}}(X \boxtimes Y, gX \boxtimes hY) \cong \mathrm{Hom}_{\mathcal{V}}(X, gX) \otimes_k \mathrm{Hom}_{\mathcal{W}}(Y, hY)$$

(see [GK, Sec.1]). Under this identification,

$$\mu\left(\sum_i \phi_i \otimes \psi_i\right)_{(X,Y)} = \sum_i \phi_{i,X} \otimes \psi_{i,Y}.$$

To define an inverse of μ, let

$$\eta \colon \boxtimes \Rightarrow \boxtimes \circ (g, h)$$

be a natural transformation. Fix an object Y of $\mathcal{W}$, and choose a basis $B^Y = (\beta_1^Y, \dots, \beta_m^Y)$ of $\mathrm{Hom}_{\mathcal{W}}(Y, hY)$. For each $X \in \mathrm{ob}\mathcal{V}$, there is a unique way to express $\eta_{(X,Y)}$ in the form

$$\eta_{(X,Y)} =: \sum_{i=1}^{m} \phi_i^X \otimes \beta_i^Y.$$

As X varies, naturality of η implies that, for each i, the maps ϕ_i^X form a natural transformation ϕ_i from $\mathrm{id}_{\mathcal{V}}$ to g. So,

$$\eta_Y := \sum_{i=1}^{m} \phi_i \otimes \beta_i^Y$$

is a well-defined element of

$$\mathbb{T}r_{\mathcal{V}}(g) \otimes \mathrm{Hom}_{\mathcal{W}}(Y, hY).$$

Without loss of generality, we assume $\mathbb{T}r(g)$ to be finite dimensional. Then we may pick a k-basis of $\mathbb{T}r_{\mathcal{V}}(g)$ and repeat our argument. This yields an element

$$\nu(\eta) \in \mathbb{T}r_{\mathcal{V}}(g) \otimes_k \mathbb{T}r_{\mathcal{W}}(h).$$

By construction, ν is a right-inverse to μ. To see that ν is also a left-inverse, we pick $\phi \in \mathbb{T}r(g)$ and $\psi \in \mathbb{T}r(h)$ and calculate $\nu(\mu(\phi \otimes \psi))$. Fix Y, and write ψ_Y as a linear combination of the basis B^Y,

$$\psi_Y =: \sum_i a_i \beta_i^Y.$$

In the construction of ν, this implies $\phi_i^X = a_i \phi_X$, hence $\phi_i = a_i \phi$ and

$$\begin{aligned} \eta_Y &= \sum_i \phi_i \otimes \beta_i \\ &= \sum_i a_i \phi \otimes \beta_i \\ &= \phi \otimes \sum_i a_i \beta_i \\ &= \phi \otimes \psi_Y. \end{aligned}$$

Hence

$$\nu(\mu(\phi \otimes \psi)) = \phi \otimes \psi,$$

as claimed, and we have proved that ν is a left-inverse of μ. $\square$

2.2 Examples

2.2.1 2-Vector spaces

Let $\mathcal{V}$ be a semisimple linear category with finitely many isomorphism classes of simple objects. Let m be the number of isomorphism classes of simple objects in $\mathcal{V}$. Then there is an equivalence of categories

$$\mathcal{V} \simeq \mathrm{Vect}_k^m .$$

This is the reason why categories like $\mathcal{V}$ are called 2-*vector spaces* [KV94]. If $\mathcal{V}$ and $\mathcal{W}$ are 2-vector spaces, Osorno has independently found an isomorphism

$$\mathbb{T}r(g \boxtimes h) \cong \mathbb{T}r(g) \otimes \mathbb{T}r(h)$$

[Oso10]. We give a brief summary of her argument: a linear functor between two 2-vector spaces can be represented by a matrix with entries in Vect_k. Its categorical trace is the direct sum of the diagonal entries. Given equivalences $\mathcal{V} \simeq \mathrm{Vect}_k^m$ and $\mathcal{W} \simeq \mathrm{Vect}_k^n$, one obtains an equivalence

$$\mathcal{V} \boxtimes \mathcal{W} \simeq \mathrm{Vect}_k^{mn} .$$

Let g be a linear endofunctor of $\mathcal{V}$, and let h be a linear endofunctor of $\mathcal{W}$. As one would expect, the matrix for $g \boxtimes h$ is the "Kronecker product" of the matrices for g and for h. The classical computation for

$$tr(g) \cdot tr(h) = tr(g \otimes h)$$

goes through and shows that $\mathbb{T}r(g \boxtimes h)$ is isomorphic to $\mathbb{T}r(g) \otimes \mathbb{T}r(h)$.

2.2.2 Algebras

Let A be an associative and unitary finite dimensional k-algebra, and let A-mod$^{\mathrm{f}}$ be the category of finitely presented right A-modules.[4] Let g be a right-exact endofunctor of $\mathrm{A\text{-}mod^f}$. Recall (e.g. from [Del90]) that g is uniquely determined by the A-A-bimodule $M := g(A)$. Here the left module structure on M comes from the left action of A on itself:

---

[4] This is the category denoted $(A)_{\mathrm{coh}}$ in [Del90].

$$A \longrightarrow \mathrm{End}(A) \longrightarrow \mathrm{End}(M).$$

The categorical trace of g is isomorphic to the center of M:

$$\begin{aligned} \mathbb{T}r(g) &\cong \mathrm{Hom_{A\text{-}A}}(A, g(A)) \\ &= \mathrm{Center}_A(M). \end{aligned}$$

Here $\mathrm{Hom_{A\text{-}A}}$ stands for morphisms of A-A-bimodules, and the second isomorphism identifies $f \in \mathrm{Hom_{A\text{-}A}}(A, M)$ with the element $f(1)$ in M.

Let B be a second finite dimensional k-algebra. Let h be a right-exact linear endofunctor of $\mathrm{B\text{-}mod^f}$. Recall from [Del90, Prop.5.3] that we have an equivalence of abelian categories

$$\left(\mathrm{A\text{-}mod^f}\right) \widehat{\boxtimes} \left(\mathrm{B\text{-}mod^f}\right) \simeq (\mathrm{A} \otimes \mathrm{B})\text{-}\mathrm{mod^f}, \tag{4}$$

where $\widehat{\boxtimes}$ is the Deligne tensor product.

Let $M := g(A)$ and $N := h(B)$. Under the equivalence (4), the isomorphism

$$\mu\colon \mathbb{T}r(g) \otimes \mathbb{T}r(h) \longrightarrow \mathbb{T}r(g \widehat{\boxtimes} h)$$

of Theorem 2.5 is identified with the canonical map

$$\mu\colon \mathrm{Center_A}(M) \otimes \mathrm{Center_B}(N) \longrightarrow \mathrm{Center_{A\otimes B}}(M \otimes N).$$

The fact that μ is an isomorphism is the degree zero part of the Künneth theorem for Hochschild cohomology (c.f. [ML95, X.7.4]).

There is, of course, overlap between this example and the previous one: if A is semisimple, then $\mathrm{A\text{-}mod^f}$ is a 2-vector space. So,

$$\mathrm{A\text{-}mod^f} \simeq \mathrm{Vect_k}^r,$$

where r is the number of isomorphism classes of simple A-modules.

Example 2.6. Take $g = \mathrm{id}$ to be the identity functor. We have seen two different ways to calculate the categorical trace of id: on one hand,

$$\mathbb{T}r(\mathrm{id}) \cong \mathrm{Center}(A).$$

On the other hand, we may calculate $\mathbb{T}r(\mathrm{id})$ as the direct sum of the diagonal entries of the $r \times r$ identity 2-matrix,

$$\mathbb{T}r(\mathrm{id}) \cong k^{\oplus r}.$$

So, we have recovered the well known fact that

$$\dim(\mathrm{Center}(A)) = r.$$

More generally, let M be an A-A-bimodule. Then the same argument yields

$$\dim(\mathrm{Center}_{\mathrm{A}}(M)) = \sum m_i,$$

where m_i is the number of times that the simple object W_i occurs as summand inside $W_i \otimes M$. Here the sum runs over all isomorphism classes of simple objects of A-$\mathrm{mod}^{\mathrm{f}}$.

2.2.3 Coherent sheaves

Let X be a smooth projective variety over k, and let $\mathcal{C}\mathrm{oh}(X)$ be the category of coherent sheaves on X. Let $\mathcal{D}^{+}_{\mathrm{coh}}(X)$ be the derived category of bounded below complexes in $\mathcal{C}\mathrm{oh}(X)$. This is a triangulated category, and it possesses an enhancement $\mathcal{I}(X)$ such that, for any two varieties X and Y as above, the completed tensor product

$$\mathcal{I}(X) \,\widehat{\boxtimes}\, \mathcal{I}(Y)$$

is quasi-equivalent to $\mathcal{I}(X \times Y)$ (see [BLL04, Th. 5.5] compare also [GK, (1.7.4)(c)]). Let g be an automorphism of X, and let h be an automorphism of Y. Then we have the direct image functors g_* and h_* acting on, respectively, $\mathcal{I}(X)$ and $\mathcal{I}(Y)$. The above equivalence identifies $g_* \widehat{\boxtimes} h_*$ with $(g \times h)_*$. Applying Theorem 2.5, we conclude

$$\mathbb{T}r^{\bullet}(g_*) \otimes \mathbb{T}r^{\bullet}(h_*) \cong \mathbb{T}r^{\bullet}((g \times h)_*).$$

These categorical traces in $\mathcal{I}(X)$ are hard to get a handle on. Instead, one often works with bicategories, compare to Section 3.4.

3 Traces in monoidal 2-categories

By a *monoidal 2-category* we will mean a k-linear semistrict monoidal 2-category in the sense of [BN96]. In this setup, the axioms still imply the existence of $g \boxtimes h$, unique up to a specified isomorphism and the existence of a map

$$\mu\colon \mathbb{T}r(g) \otimes \mathbb{T}r(h) \to \mathbb{T}r(g \boxtimes h).$$

In general, there is no reason for μ to be an isomorphism, but in the following we will see some examples where it is. To be precise, the lax 2-categories in the following sections would need to be strictified in an appropriate sense (see [KV94, 2.12, 4.3]) in order for the formalism of [BN96] to apply. Rather than working out the precise formalism in the lax case, we choose an ad hoc approach to the individual examples.

3.1 Bimodules, strict version

We recall from [GK08, 2.2(c)] the k-linear 2-category $\mathcal{B}\mathrm{im}_k$. The objects of $\mathcal{B}\mathrm{im}_k$ are associative and unitary k-algebras. Given two such algebras, A and B, the category of 1-morphisms from A to B

$$1\,\mathrm{Hom}_{\mathcal{B}\mathrm{im}_\mathrm{k}}(A, B) = \mathcal{B}\mathrm{im}_\mathrm{k}(A, B)$$

is the category of A-B-bimodules (where the left and right actions of k are required to agree). Horizontal composition of 1-morphisms into and out of A is given by the tensor product over A. Again, the degree zero part of the Künneth theorem for Hochschild cohomology (c.f. [ML95, X.7.4]) gives the isomorphism

$$\begin{aligned}\mu\colon \mathbb{T}r(M) \otimes \mathbb{T}r(N) &\cong \mathrm{Center}_\mathrm{A}(M) \otimes \mathrm{Center}_\mathrm{A}(N)\\ &\cong \mathrm{Center}_\mathrm{A}(M \otimes N)\\ &\cong \mathbb{T}r(M \otimes N).\end{aligned}$$

This is closely related to Section 2.2.2: Let A and B be associative and unitary finite dimensional k-algebras. Let $\mathcal{B}\mathrm{im}^\mathrm{f}_\mathrm{k}(A, B)$ be the full subcategory of $\mathcal{B}\mathrm{im}_\mathrm{k}(A, B)$ whose objects are bimodules that are finitely presented as right B-modules. Then there is an equivalence of categories

$$\begin{aligned}\mathcal{F}\mathrm{un}_{\mathrm{r.e.}}\left(\mathrm{A\text{-}mod}^\mathrm{f}, \mathrm{B\text{-}mod}^\mathrm{f}\right) &\simeq \mathcal{B}\mathrm{im}^\mathrm{f}_\mathrm{k}(A, B) \qquad (5)\\ F &\longmapsto F(A)\\ (-) \otimes_A M &\longleftarrow M.\end{aligned}$$

Here $\mathcal{F}\mathrm{un}_{\mathrm{r.e.}}$ stands for the category of k-linear right-exact functors and natural transformations between them. Under this equivalence, composition of functors corresponds to composition of 1-morphisms in $\mathcal{B}\mathrm{im}_k$.

3.2 Bimodules, derived version

The triangulated k-linear 2-category $\mathcal{D}\mathcal{B}\mathrm{im}_k$ of [GK08, 2.4(b), 3.5] has the same objects as $\mathcal{B}\mathrm{im}_k$. The category

$$1\mathrm{Hom}_{\mathcal{D}\mathcal{B}\mathrm{im}_k}(A,B)$$

is the derived category of complexes of A-B-bimodules bounded above. Horizontal composition is as above, but using the derived tensor product. Let A be a finite dimensional, associative and unitary k-algebra, viewed as an object of $\mathcal{D}\mathcal{B}\mathrm{im}_k$. Let $K^\bullet$ be a 1-endomorphism of A. Then the categorical trace of $K^\bullet$ is the hypercohomology of $K^\bullet$,

$$\begin{aligned} \mathbb{T}r^\bullet(K^\bullet) &= \mathrm{Hom}^\bullet_{\mathcal{D}^+\mathcal{B}\mathrm{im}(A-A)}(A,K^\bullet) \\ &= \mathbb{H}^\bullet(A;K^\bullet). \end{aligned}$$

In the special case were $K^\bullet$ is a single A-A-bimodule M situated in degree zero,

$$\begin{aligned} \mathbb{T}r^\bullet(M) &= \mathrm{Ext}^\bullet_{A\otimes A^{\mathrm{op}}}(A,M) \\ &= \mathrm{HH}^\bullet(A,M) \end{aligned}$$

is the Hochschild cohomology of M. Let B be another finite dimensional, associative and unitary k-algebra, and let $L^\bullet$ be a complex of B-B-bimodules, bounded above. Then

$$\mu\colon \mathbb{T}r^\bullet(K^\bullet)\otimes\mathbb{T}r^\bullet(L^\bullet)\longrightarrow\mathbb{T}r(K^\bullet\otimes L^\bullet)$$

is the Künneth map. In the special case where both $K^\bullet$ and $L^\bullet$ are single bimodules situated in degree zero, the Künneth theorem for Hochschild cohomology (c.f. [ML95, X.7.4]) implies that μ is an isomorphism.

3.3 Constructible sheaves

The 2-category $\mathbb{R}\mathcal{A}\mathrm{n}_{\mathbb{C}}$ of [GK08, 2.4.(d)] has as objects real analytic manifolds. For any two such manifolds X and Y, the category of 1-morphisms between them is defined as

$$1\mathrm{Hom}_{\mathbb{R}\mathcal{A}\mathrm{n}_{\mathbb{C}}}(X,Y) = \mathcal{D}^{+}_{\mathrm{constr}}(X \times Y),$$

the bounded derived category of ($\mathbb{R}$-)constructible sheaves of $\mathbb{C}$-vector spaces on $X \times Y$.[5] Horizontal composition of

$$\mathcal{K} \in 1\mathrm{Hom}(X,Y) \quad \text{and} \quad \mathcal{L} \in 1\mathrm{Hom}(Y,Z)$$

is given by the *derived convolution*

$$\mathcal{L} \circ \mathcal{K} = R\pi_{13*}\left(\pi_{12}^{-1}\mathcal{K} \otimes^{L} \pi_{23}^{-1}\mathcal{L}\right).$$

Here the π_{ij} are the projections from $X \times Y \times Z$ to the according products with two factors, the tensor product is taken over the constant sheaf $\underline{\mathbb{C}}$, and the π_{ij}^{-1} are the respective inverse image sheaf functors.

Every object ("kernel") $\mathcal{K} \in 1\mathrm{Hom}(X,Y)$ defines a functor

$$\begin{array}{rcl} F_{\mathcal{K}}\colon \mathcal{D}^{+}_{\mathrm{constr}}(X) & \longrightarrow & \mathcal{D}^{+}_{\mathrm{constr}}(Y) \\ \mathcal{F} & \longmapsto & R\pi_{Y*}\left(\pi_X^{-1}\mathcal{F} \otimes^{L} \mathcal{K}\right). \end{array}$$

Let $g\colon X \to X$ be analytic, and let

$$\gamma\colon X \to X \times X$$

be the inclusion of the graph Γ_g. Consider the kernel

$$\mathcal{K}_g = \underline{\mathbb{C}}_{\Gamma_g} := \gamma_* \underline{\mathbb{C}},$$

the constant sheaf supported on Γ_g, viewed as a complex situated in degree 0. This is a lift of g_* to $\mathbb{R}\mathcal{A}\mathrm{n}_{\mathbb{C}}$. In other words, we have $F_{\mathcal{K}_g} = g_*$. We have[6]

$$\mathbb{T}r^{\bullet}(\mathcal{K}_g) \cong \bigoplus_{\alpha} H^{\bullet - codim X^g_\alpha}(X^g_\alpha; \underline{\mathbb{C}}),$$

where the sum runs over the connected components of the fixed point set. Let Y be a second object of $\mathbb{R}\mathcal{A}\mathrm{n}_{\mathbb{C}}$, and consider an analytic automorphism h of Y. Then the connected component $X^g_\alpha \times Y^h_\beta$ of $(X \times Y)^{(g,h)}$ has codimension

$$\mathrm{codim}(X^g_\alpha) + \mathrm{codim}(Y^h_\beta) \tag{6}$$

---

[5] See [KS94, Section 8.4], for background on constructible sheaves.

[6] This is Proposition 5.5 in [GK08]. Note the sign mistake there: in the proof of Proposition 5.5, $\underline{\mathbb{C}}_{X^g}[\mathrm{codim}(X^g)]$ should be $\oplus \underline{\mathbb{C}}_{X^g_\alpha}[-\mathrm{codim}(X^g_\alpha)]$.

in $X \times Y$. The map

$$\mu\colon H^\bullet(X^g, \underline{\mathbb{C}}) \otimes H^\bullet\left(Y^h, \underline{\mathbb{C}}\right) \longrightarrow H^\bullet\left(X^g \times Y^h, \underline{\mathbb{C}}\right)$$

is the Künneth isomorphism. Here the notation is short-hand for the direct sum over the connected components of $X^g \times Y^h$, with the grading in each summand shifted by (6).

3.4 Coherent sheaves

Another key example is the 2-category $\mathcal{V}\mathrm{ar}_{\mathbb{C}}$.[7] Its objects are smooth projective varieties over $\mathbb{C}$, and for two such varieties X and Y, the category of 1-morphisms from X to Y is

$$1\mathrm{Hom}(X, Y) = \mathcal{D}^+_{\mathrm{coh}}(X \times Y),$$

the derived category of coherent sheaves on $X \times Y$. The formalism is identical to that of the previous section, but now the derived tensor product $\otimes^L$ is taken over the structure sheaf, and the inverse image functors π_{ij}^{-1} are replaced by the pull-back functors π_{ij}^*. For an automorphism g of X, the kernel

$$\mathcal{K}_g = \mathscr{O}_{\Gamma_g} := \gamma_* \mathscr{O}_X$$

lifts the auto-equivalence g_* of $\mathcal{D}^+_{\mathrm{coh}}(X)$ in the sense explained in the previous section. Its trace is identified by the following theorem:

Theorem 3.1. *Let N^g be the normal bundle of the inclusion*

$$u\colon X^g \hookrightarrow X.$$

Then we have an isomorphism

$$\mathbb{T}r^\bullet(g) \cong \bigoplus_\alpha \mathrm{HH}^{\bullet\text{-codim}(\mathrm{X}^{\mathrm{g}}_\alpha)}\left(X^g_\alpha, \det(N^g)\right),$$

where the direct sum is over the connected components of the fixed point set X^g.

---

[7] This is [GK08, 2.4.Exa(c)]. For a detailed account, see [CW10] or [Huy06].

Again,

$$\mu\colon H^\bullet(X^g, \det(N^g)) \otimes H^\bullet\left(Y^h, \det(N^h)\right) \longrightarrow H^\bullet\left(X^g \times Y^h, \det(N^{(g,h)})\right)$$

is the Künneth isomorphism. Again, the notation is short-hand for the direct sum over the connected components of $X^g \times Y^h$, with the grading in each summand shifted by (6).

Proof of Theorem 3.1: We have a commuting diagram of closed immersions

$$\begin{tikzcd}
X^g \arrow[rr, "u"] \arrow[dd, "u"'] \arrow[dr, "\delta"] & & X \arrow[dd, "\gamma"] \\
& X^g \times X^g \arrow[dr, "i"] & \\
X \arrow[rr, "\Delta"'] & & X \times X.
\end{tikzcd} \tag{7}$$

Here Δ and δ denote the inclusions of the respective diagonals, and γ is the inclusion of the graph Γ_g. All squares in (7) are cartesian. We have

$$\begin{aligned}
\mathbb{T}r^\bullet(g) &= \mathrm{Ext}^\bullet_{\mathrm{X\times X}}\left(\Delta_*\mathscr{O}_X, \gamma_*\mathscr{O}_X\right) \\
&= \mathrm{Ext}^\bullet_{\mathrm{X}}\left(L\gamma^*\Delta_*\mathscr{O}_X, \mathscr{O}_X\right).
\end{aligned}$$

Write Θ for the composite of natural transformations

$$\Theta\colon L\gamma^*\Delta_* \Longrightarrow L\gamma^*\Delta_* u_* Lu^* = L\gamma^* i_*\delta_* Lu^* \Longrightarrow u_* L\delta^*\delta_* Lu^*,$$

where the first map is given by the unit of the adjunction $Lu^* \dashv u_*$, and the second map is given by the base-change map of the upper right (triangle shaped) square in (7).



The natural transformation Θ in string diagram notation. The squares labeled σ and τ both denote the identity natural transformation of $(\gamma u)_* = (i\delta)_* = (\Delta u)_*$.

We claim that Θ is an isomorphism. The claim implies the theorem, by the following calculation:

$$\begin{aligned}\mathrm{Ext}^{\bullet}_{\mathrm{X}}(L\gamma^{*}\Delta_{*}\mathscr{O}_X,\mathscr{O}_X) &\cong \mathrm{Ext}^{\bullet}_{\mathrm{X}}(u_{*}L\delta^{*}\delta_{*}\mathscr{O}_{X^g},\mathscr{O}_X)\\ &\cong \mathrm{Ext}^{\bullet}_{\mathrm{X^g\times X^g}}(\delta_{*}\mathscr{O}_{X^g},\delta_{*}u^{!}\mathscr{O}_X)\\ &= \mathrm{HH}^{\bullet}\big(X^g,u^{!}\mathscr{O}_X\big).\end{aligned}$$

Here $u^!$ is the right-adjoint of u. Explicitly, $u^!$ is given by

$$u^{!}\mathscr{O}_X \cong \bigoplus_{a}\det(N^g_\alpha)[-\operatorname{codim}(X^g_\alpha)]$$

(see e.g. [Căl05].)

The proof that Θ is an isomorphism is an adaptation of the discussion in [CKS03, Appendix A]. As the question is local, it will suffice to prove it over an open cover of $X\times X$. For $(x,y)\in X\times X\smallsetminus\Delta(X^g)$, there exists an open neighourhood U of (x,y) with $U\cap\Delta=\varnothing$ or $U\cap\Gamma_g=\varnothing$. Since $\varnothing=\operatorname{spec}(\{0\})$, the claim is trivially true over such a U. Let now $x\in X^g$. Writing

$$T_{\Delta(X)}:=\operatorname{im}(T_\Delta),\quad T_{\Gamma_g}=\operatorname{im}(T_\gamma),\quad\text{and}\quad T_{\Delta(X^g)}:=\operatorname{im}(T_{i\delta}),$$

we have

$$T_{\Delta(X)}\cap T_{X^g\times X^g}=T_{\Delta(X^g)}=T_{\Delta(X)}\cap T_{\Gamma_g}$$

(inside $T_{X\times X}$). By Lemma A.5, we have

$$T_{\Delta(X)\cup(X^g\times X^g)}=T_{\Delta(X)}+T_{X^g\times X^g}.$$

We now apply Lemma A.3 with $Z=X\times X$ and $W=\Delta(X)\cup(X^g\times X^g)$. Over a neighbourhood U of x inside $X\times X$ this yields a refinement of (7) looking as follows:

$$\begin{tikzcd}
X^g \arrow[rr, "u"] \arrow[dd, "u"'] \arrow[dr, "\delta"] & & X \arrow[dd, "\gamma"] \\
& X^g\times X^g \arrow[d, "j"'] \arrow[dr, dotted, "i"] & \\
X \arrow[r, "\Delta'"'] & Y \arrow[r, "k"'] & X\times X
\end{tikzcd}$$

(all objects intersected with U). Here Y is a smooth subvariety of $X\times X$, the maps j, k, and Δ' are closed immersions, and

$$T_{Y,x}=T_{\Delta(X),x}+T_{X^g\times X^g,x}.$$

The left solid square (with arrows δ, j u and Δ') is still cartesian. Further, $T_{i\delta}$ defines an isomorphism

$$T_{X^g,x} \cong T_{T,x} \cap T_{\Gamma_g,x}$$

(intersection inside $T_{X\times X,x}$). Applying Lemma A.6, we conclude that (possibly after passing to a smaller neighbourhood of x) the right solid square (with arrows u, γ,, $j\delta$ and k) is also cartesian. Moreover, both intersections

$$X^g = \Delta(X) \cap_Y (X^g \times X^g) \quad \text{and} \quad X^g = Y \cap_{X\times X} \Gamma_g$$

are clean and "of the expected dimension", meaning that

$$\dim(X^g) + \dim(Y) = \dim(X) + \dim(X^g \times X^g)$$

and similarly for $Y \cap \Gamma_g$. Using [Ser77, p.V 20, Cor. to Theorem 4], we conclude that both these squares are *tor-indepent squares* in the sense of [Lip09, Def. (3.10.2)]. By [Lip09, Thm. (3.10.3)], the base change tranformations

$$L\gamma^* k_* \Longrightarrow u_* L(j\delta)^*$$

and

$$L_j^* \Delta'_* \Longrightarrow \delta_* L u^*$$

are isomorphisms. Hence, over U we have a natural isomorphism

$$\Theta' \colon L\gamma^* \Delta_* \Longrightarrow u_* L\delta^* Lj^* \Delta'_* \Longrightarrow u_8 L\delta^* \delta_* L u^*.$$



The natural transformation Θ' in string diagram notation. The squares labeled σ and τ' denote the identity natural transformations of $(\gamma u)_* = (kj\delta)_*$ and of $(\Delta' u)_* = (j\delta)_*$. The part differing from the diagram for Θ is red.

To see that $\Theta' = \Theta|_U$, we move τ' downward until it is below σ, then straighten the red zig-zag line connecting σ and τ' into a straight line labeled j_*, and finally note that

$$k_*\tau' = \mathrm{id}_{(i\delta)_*} = \tau.$$

□

4 Tensor products of 2-representations

Let G be a finite group. Recall, e.g. from [GK08, Def.4.1.], that a 2-representation of G on a k-linear category $\mathcal{V}$ consists of the following data:

- for each $g \in G$ an endofunctor $\varrho(g)$ of $\mathcal{V}$,
- for any $g, h \in G$, a natural isomorphism

  $$\phi_{g,h} \colon \varrho(g)\varrho(h) \xrightarrow{\cong} \varrho(gh),$$

  and

- a natural isomorphism

$$\phi_1\colon \varrho(1) \overset{\cong}{\Longrightarrow} \mathrm{id}_{\mathcal{V}},$$

The $\phi_?$ are required to make associativity pentagons and unit diagrams commute. In particular, for g, h and k in G we have an unambiguous double composition isomorphism

$$\phi_{g,h,k}\colon \varrho(g)\varrho(h)\varrho(k) \overset{\cong}{\Longrightarrow} \varrho(ghk).$$

Following Bartlett[8], we use the string diagram notation and write



for $\phi_{g,h}$,



for $\phi_{g,h}^{-1}$,



for ϕ_1,



for ϕ_1^{-1},



for $\phi_{g,h,k}$,



for $\phi_{g,h,k}^{-1}$.

We recall further that $\varrho(g^{-1})$ is a weak inverse, and hence a two-sided adjoint, of $\varrho(g)$. For the adjunction $\varrho(g^{-1}) \dashv \varrho(g)$, we have



The unit is given by $(\phi_{g,g^{-1}}\phi_1)^{-1}$ and denoted by either of these two diagrams,



and the counit is given by $\phi_{g^{-1},g}\phi_1$ and denoted by either of these two diagrams.

The axioms for the $\phi_{g,h}$ imply that these adjunctions compose as one would hope for, i.e., the adjunction for g composed with that for s gives the adjunction for sg. We refer the reader to [Bar08], in particular Lemma 6 (iv), for a detailed account of these facts and their expression in terms of string diagram moves.

---

[8] See [Bar08, p.13] for a translation of the axioms into string diagram moves. Note that our string diagrams flow upwards, while Bartlett's flow downwards.

The categorical character of ϱ consists of the data

$$X_\varrho(g) = \mathbb{T}r(\varrho(g)), \quad g \in G$$

and for each pair (s,g) the isomorphism

$$\psi_s \colon X_\varrho(g) \longrightarrow X_\varrho(sgs^{-1}),$$

sending $\xi \in \mathbb{T}r(\varrho(g))$ to



$\psi_s(\xi) \in \mathbb{T}r(\varrho(sgs^{-1}))$

and satisfying $\psi_1 = \mathrm{id}$ and $\psi_s\psi_t = \psi_{st}$ (see [GK08, Prop.4.10] or [Bar08, Prop.16] for details).

Definition 4.1. Let ϱ be a 2-representation of G on a linear category $\mathcal{V}$, and let π be a 2-represenation of H on a linear category $\mathcal{W}$. We define the 2-representation $\varrho \boxtimes \pi$ on $\mathcal{V} \boxtimes \mathcal{W}$ by applying $- \boxtimes -$ to all the data of 2-representation.

Corollary 4.2. *In the situation of Theorem 2.5, we have isomorphisms*

$$\mu \colon X_\varrho(g) \otimes X_\pi(h) \xrightarrow{\cong} X_{\varrho\boxtimes\pi}(g,h)$$

identifying $\psi_s^\varrho \otimes \psi_t^\pi$ *with* $\psi_{(s,t)}^{\varrho\boxtimes\pi}$. *In orther words,* $X_\varrho \otimes X_\pi$ *and* $X_{\varrho\boxtimes\pi}$ *are isomorphic as representations of the inertia groupoid* $\Lambda(G \times H)$.

Proof : This follows from Theorem 2.5 and inspection of the definition of ψ_s in the proof of [GK08, Prop.4.10.]. ☐

A similar statement holds for $\widehat{\boxtimes}$ in the abelian case and, more generally, in any monoidal 2-category where μ is an isomorphism.

5 Categories of equivariant objects

Let $\varrho\colon G \to GL(V)$ be a complex representation of a finite group. We write V^G for the maximal G-invariant summand of V. The dimension of V^G is

$$\begin{aligned} \dim(V^G) &= \langle V, \mathbb{C}\rangle_G \\ &= \frac{1}{|G|}\sum_{g\in G}\chi_\varrho(g), \end{aligned} \tag{8}$$

where χ_ϱ is the character of ϱ.

This section studies categorical analogues of V^G. These are the category of equivariant objects and its generalizations.

5.1 Background and Definitions

The notion of *category of equivariant objects* already plays a role in [Gro57], see page 196 of [loc.cit].

Definition 5.1. Let $\mathcal{V}$ be a linear category, acted upon from the left by a finite group G. An *equivariant object* of $\mathcal{V}$ consists of an object $x \in \mathrm{ob}(\mathcal{V})$ and a system of isomorphisms

$$\{\epsilon_g\colon x \to gx\}_{g\in G}$$

such that for any $g, h \in G$ the diagram

$$\begin{tikzcd} x \arrow[r, "\epsilon_g"] \arrow[d, "\epsilon_{gh}"'] & gx \arrow[d, "g\epsilon_h"] \\ (gh)x & g(hx) \arrow[l, "\phi_{g,h,x}"] \end{tikzcd} \tag{9}$$

is commutative. As a consequence, we obtain the unit condition

$$\epsilon_1 = \phi_{1,x}^{-1} : x \to 1x.$$

A *map of equivariant objects*

$$f\colon (x,\epsilon) \to (y,\delta)$$

is a morphism $f \in \mathrm{Hom}(x,y)$ satisfing $\forall g \in G$

$$g(f)\circ\epsilon_g = \delta_g \circ f.$$

The category of equivariant objects and their maps is denoted $\mathcal{V}^G$.

Let X be a G-space. Then the categories of equivariant k-vector bundles over X, equivariant sheaves over X, and equivariant spaces over X have interpretations as categories of equivariant objects. Below, we will discuss some of these examples in detail. Maps of 2-representations form another example:

Definition 5.2. Let $\mathcal{C}$ be a 2-category, and let π and ϱ be 2-representations of G on objects W and V of $\mathcal{C}$. Then $G \times G$ acts on the category $1\mathrm{Hom}(W, V)$ by

$$F \longmapsto \varrho(g) F \pi(s^{-1}).$$

The coherence isomorphisms of this action are inherited from those of π and ϱ.

View G as a subgroup of $G \times G$ via the diagonal inclusion. We write

$$1\mathrm{Hom}_{\mathrm{G}}(W, V) := 1\mathrm{Hom}(W, V)^{G},$$

for the *category of equivariant 1-homohorphisms and G-invariant 2-homomorphisms between them*; and

$$1\mathrm{Hom}_{\mathrm{G}}^{\mathrm{triv}}(W, V) := 1\mathrm{Hom}(W, V)^{G \times G}$$

for the category of *trivialized G-equivariant 1-morphisms and trivialization-preserving G-invariant 2-homomorphisms.* Explicitly, an object of $1\mathrm{Hom}_{\mathrm{G}}(W, V)$ consists of a 1-morphism $f \colon W \to V$ together with a compatible family of 2-isomorphisms

$$\eta_g \colon F \Longrightarrow \varrho(g) F \pi(g).$$

An alternative point of view is to consider the flip 2-isomorphisms τ_g given by the string diagrams



Definition of τ_g



Notation for τ_g



Notation for τ_g^{-1}

and satisfying

$$\tau_1 = \phi_1 F \phi_1^{-1}$$

(horizontal composition but vertical inverse) and

$$\tau_{gh} \circ (\phi_{g,h} F) = (F\phi_{g,h}) \circ (\tau_g h) \circ (g\tau_h)$$

(see [Bar08, (5)]). The data $(F, \{\tau_g\})$ are eqivalent to those of $(F, \{\eta_g\})$. An object of $1\mathrm{Hom}_{\mathrm{G}}^{\mathrm{triv}}$ can then be thought of as a G-equivariant 1-morphism $(F, \{\tau_g\})$ from V to W together with a family of *(left) trivilization 2-isomorphisms*

$$\epsilon_g \colon F \Longrightarrow \varrho(g)F,$$

satisfying (9) and compatible with $\{\tau_s\}$ in the following sense:



Notation for ϵ_g



These are equal to each other.

Equivalently, one may work with the right trivialization isomorphisms

$$\epsilon_g^R := \tau_g \circ \epsilon_g$$

with symmetric notation and compatibility requirements to those of ϵ_g (here $\eta_{(g,1)}$ is replaced by $\eta_{(1,g^{-1})}$). In the special case where $F = \mathrm{id}_W$, we say that the G-action on W is trivialized (by $\{\epsilon_g\}$).

Example 5.3. Let $F \colon V \to W$ and $H \colon W \to Z$ be equivariant 1-morphisms, write $\{\tau_g\}$ and $\{\sigma_g\}$ for the respective families of twist maps. Then the composition HF, together with the twist maps $(H\tau_g) \circ (\sigma_g F)$, is again an equivariant 1-morphism. The horizontal composition of G-invariant 2-morphisms remains G-invariant.

If H is trivialized by $\{\epsilon_s\}$ then HF inherits the trivialization maps $\epsilon_g F$. If F is trivialized with right trivialization maps $\{\delta_s^R\}$ then HF inherits a trivialization with right trivialization maps $\{H\delta_s^R\}$. If both H and F are trivialized then the inherited trivilizations of HF might not agree.

Example 5.4. The identity 1-automorphism of V is equivariant, with the η_g given by the units of the adjunctions $g^{-1} \dashv g$. The fact that these η_g compose in the expected way is (9).

If there exists a family $\{\epsilon_g : \mathrm{id}_V \Rightarrow \varrho(g)\}$ satisfying (9) then η_g is identified with the horizontal composition

$$\eta_g = \epsilon_g \epsilon_{g^{-1}}. \tag{10}$$

It follows that $\{\epsilon_g\}$ is automatically compatible with $\{\tau_g\}$ So, ϱ is trivialized by $\{\epsilon_g\}$. Further, $\epsilon_g^R = \epsilon_g$.

In this situation, we have fully faithful embeddings

$$\begin{array}{rcl} 1\mathrm{End}(V) & \longrightarrow & 1\mathrm{End}_{\mathrm{G}}(V) \\ F & \longmapsto & \big(F, \{\epsilon_g F \epsilon_{g^{-1}}\}\big). \end{array} \tag{11}$$

and

$$\begin{array}{rcl} 1\mathrm{Hom}_{\mathrm{G}}(V, W) & \longrightarrow & 1\mathrm{Hom}_{\mathrm{G}}^{\mathrm{triv}}(V, W) \\ H & \longmapsto & (H, \{H\epsilon_g\}) \end{array} \tag{12}$$

for any other object W with G-action.

We are now ready to formulate the universal property characterizing $\mathcal{V}^G$.

Proposition 5.5. *There is a G-action on $\mathcal{V}^G$, which is canonically trivialized by the structure maps $\{\epsilon_g\}$. The forgetful functor*

$$U \colon \mathcal{V}^G \longrightarrow \mathcal{V}$$

is G-equivariant. Assume we are given another category $\mathcal{W}$ with a G-action. Then composition with U defines an equivalence of categories

$$\mathcal{F}\mathrm{un}_G(\mathcal{W}, \mathcal{V}^G) \xrightarrow{\simeq} \mathcal{F}\mathrm{un}_G^{triv}(\mathcal{W}, \mathcal{V}).$$

Here the trivialization of UF is inherited from the trivialization $\{\epsilon_g\}$ of $\mathrm{id}_{\mathcal{V}^G}$ (see Example 5.3). In pictures:



These two diagrams agree.



These two diagrams agree.



These two diagrams agree.

Proof : On objects of $\mathcal{V}^G$, the G-action is defined by

$$s\colon \{x, \epsilon_g\} \longmapsto \{sx, s\epsilon_{s^{-1}gs}\}.$$

All the other data of G-action are inherited from the action on $\mathcal{V}$. Note that we are forced to define $\epsilon_g s$ in this manner in order for ϵ_g to be a natural transformation from id to g in $\mathcal{V}^G$:

$$(\epsilon_g s) \circ \epsilon_s = (g\epsilon_s) \circ \epsilon_g$$

is equivalent to

$$\epsilon_g s = \epsilon_{gs}\epsilon_s^{-1} = s\epsilon_{s^{-1}gs}. \tag{13}$$

By construction, the family $\{\epsilon_s\}$ satisfies the coherence conditions (9). One checks that the functor of the proposition is a well-defined equivalence of categories. □

The universal property in Proposition 5.5 should be compared that in [GK08, Prop. 4.4]. Although the latter looks less general, it is equivalent to the one here: given a trivialized functor $F\colon \mathcal{W} \to \mathcal{V}$, we can apply [GK08, Prop. 4.4] to the essential image of F and obtain the result of Proposition 5.5. This argument does, however, not go through for representations in general 2-categories, where we will work with the universal property formulated here.

Definition 5.6. Let $\mathcal{C}$ be a 2-category, and let ϱ be a 2-representation of G on $V \in \mathrm{ob}(\mathcal{C})$. We will write

$$U\colon V^G \longrightarrow V$$

whenever V^G is an object of $\mathcal{C}$ and U is an equivariant 1-morphism from V^G to V such that (V^G, U) satisfies the universal property of Proposition 5.5.

If it exists, the object V^G is well defined up to trivialization preserving equivariant 1-equivalence, which in turn is unique up to canonical, trivialization preserving, G-invariant 2-morphism. From now on, we fix $U\colon V^G \longrightarrow V$ as in Definition 5.6. We write $\{\upsilon_g\}$ for the (left) trivialization maps of the tivialization that U inherits from id_{V^G}, i.e., υ_g is the 2morphism $U\epsilon_g$ precomposed with the flip map $gU \Rightarrow Ug$.

We will often supress the representation ϱ from the notation and write g instead of $\varrho(g)$.

Proposition 5.7. *The 1-morphism U possesses a left-adjoint*

$$A' \in 1\,\mathrm{Hom}(V, V^G)$$

satisfying

$$UA' = A := \bigoplus_{g \in G} g.$$

Proof : The structure isomorphisms

$$\phi_{g,h,s^{-1}} \colon \varrho(g)\varrho(h)\varrho(s^{-1}) \xrightarrow{\cong} \varrho(ghs^{-1}).$$

make A into a trivialized equivariant 1-morphism. Write $\varphi_{s,A} \colon A \Rightarrow sA$ for the trivialization 2-morphisms. Applying the universal property of V^G, we obtain a G-equivariant 1-morphism

$$A' \colon V \longrightarrow V^G$$

with $UA' = A$. Let ι be the composite

$$\iota \colon \mathrm{id}_V \xRightarrow{\phi_1^{-1}} 1 \Longrightarrow A$$

where the second map is the inclusion of the 1st summand. let

$$\alpha \colon AU \Longrightarrow U$$

be the 2-morphism

$$\alpha := \sum_{g \in G} \upsilon_g^{-1}.$$

By (9), we have

$$\alpha \circ (\phi_{s,A} U) = \upsilon_s \circ \alpha.$$

in other words, α is a trivialization preserving (equivariant) 2-morphism

$$(AU, \{\varphi_{s,A} U\}) \Longrightarrow (U, \{\upsilon_s\}).$$

The universal property of V^G yields a 2-morphism

$$\alpha' \colon A'U \Longrightarrow 1$$

with $U\alpha' = \alpha$. We need to show that ι and α' form the unit and counit of an adjunction $A' \dashv U$. The identity

$$(U\alpha') \circ (\iota U) = \mathrm{id}_U$$

is immediate from the definitions and from $\epsilon_1 = \phi_1^{-1}$. By the universal property, it sufficies to check the second condition after horizontal composition with U. So, we are left to show

$$(\alpha A) \circ (A\iota) = \mathrm{id}_A \,.$$

Restricted to the gth summand, this composite is the left-most in the following sequence of string diagrams:









Here ι_g denotes the inclusion of the gth summand. The equality of the first three diagrams completes the proof. Their equality to the fourth will be used below. □

5.2 The twisted group algebra

Let G act on a linear category $\mathcal{V}$, and let $\mathcal{V}^G$ be the category of equivariant objects. We will think of this category as resembling the category of representations of G "in a twisted sense". Representations of G are the same as modules over the group algebra $k[G]$, and it is natural to ask whether there is an analog of $k[G]$ for this situation – an associative algebra acting on each equivariant object. The answer is: yes, and very simple.

Let ϱ be a representation of G on an object V of a linear 2-category, and let A be as in Proposition 5.7. Then, by general nonsense about adjunctions, $A = UA'$ is a monoid with multiplication

$$\alpha A' \colon A^2 \Longrightarrow A$$

and unit ι. Let F be a 1-endomorphism of V^G. Then $M := UFA'$ is a bimodule over A with the left- and right module structure given by , respecitively, $\alpha F A'$ and $UF\alpha' A'$.

Definition 5.8. The *twisted group algebra* of ϱ is the space

$$R_\varrho := \mathbb{T}r(A) = \bigoplus_{g\in G} \mathbb{T}r(g),$$

together with the k-algebra structure induced by the monoid structure of A.



The product of elements ξ and ζ of $\mathbb{T}r(A)$

If F is a 1-endomorphism of V^G, and $M = UFA'$ then $\mathbb{T}r(M)$ is a bimodule over R_ϱ.

Lemma 5.9. *Let ξ be in the summand $\mathbb{T}r(g)$ of R_ϱ, and let $\mu \in \mathbb{T}r(M)$. Then their products are expressed by the following string diagrams:*



Left multiplication



Right multiplication

Proof : The diagram for the left multiplication is just the definition of α. The diagram for the right multiplication follows from the equality of the third and the fourth diagram in the proof of Proposition 5.7. ☐

If F is of the form $\widetilde{H}$ for an equivariant 1-endomorphism $(H, \{\tau_g\})$ of V then $M = HA$, and

$$\mathbb{T}r(HA) \cong \bigoplus_{g\in G} \mathbb{T}r(Hg).$$

Proposition 5.10. *The elements $\mu \in \mathbb{T}r(Hs)$ and $\xi \in \mathbb{T}r(g)$ multiply as follows:*

$$\begin{aligned} \mu\cdot\xi &= (H\phi_{s,g})\circ(\mu\xi) \\ \xi\cdot\mu &= (H\phi_{g,s})\circ(\tau_g s)\circ(\xi\mu)\,. \end{aligned}$$

In particular, multiplication in R_ϱ is horizontal composition followed by $\phi_{g,h}$

$$\begin{array}{rcl} \mathbb{T}r(\varrho(g)) \otimes \mathbb{T}r(\varrho(h)) & \longrightarrow & \mathbb{T}r(\varrho(gh)) \\ \xi \otimes \zeta & \longmapsto & \phi_{g,h} \circ (\xi\zeta). \end{array}$$

Lemma 5.11. *There is an isomorphism of k-algebras*

$$j \colon R_\varrho \xrightarrow{\cong} 2\mathrm{End}(U),$$

where the multiplication on the target is vertical composition. For any 1-endomorphism F of V^G there is an isomorphism of R_ϱ-bimodules

$$j_F \colon 2\mathrm{Hom}(\mathrm{id}_V, UFA') \xrightarrow{\cong} 2\mathrm{Hom}(U, UF).$$

Proof : This is a consequence of the adjunction $A' \dashv U$. The isomorphism j and its inverse are defined as

$$\begin{array}{rcl} j(\xi) & = & \alpha \circ (\xi U) \\ j^{-1}(\vartheta) & = & (\vartheta A') \circ \iota, \end{array}$$

and similarly for j_F. More precisely, j_F is the second isomorphism on page 14 of [CW10] with $\Psi = A'$, $\Phi = U$, $\widehat{\Theta} = \mathrm{id}$, and $\widehat{\Xi} = UF$. □

In the situation where G acts on a linear category $\mathcal{V}$, let $(V, \{\epsilon_g\})$ be an equivariant object. Then we can compose j with the map

$$\begin{array}{rcl} 2\,\mathrm{End}(U) & \longrightarrow & \mathrm{End}_{\mathcal{V}}(V) \\ \vartheta & \longmapsto & \vartheta_V \end{array}$$

to obtain the promised action of R_ϱ on V. Explicitly, $\xi = (\xi_g)_{g \in G}$ acts on V by the endomorphism

$$\sum_g (\epsilon_g^{-1} \circ \xi_{g,V}).$$

Example 5.12. Let $\mathcal{V} = \mathrm{Vect}_k$, and let the G-action be defined by means of a 2-cocycle $c \in Z^2(G, k^*)$. So, we have the central extension

$$p \colon \widetilde{G} \to G$$

with kernel k^* and a bunch of 1-dimensional spaces

$$L_g = p^{-1}(g) \cup 0,$$

as in [GK08]. Then $\mathbb{T}r(g) = L_g$, and the algebra R_ϱ is the direct sum of all the L_g. This is the standard concept of the twisted group algebra associated to a central extension, or the cocycle. An alternative definition of this algebra is by a basis b_g, for $g \in G$ with multiplication law

$$b_g \cdot b_h = c(g,h) \cdot b_{gh}.$$

5.3 The G-action on the twisted group algebra

Let F be an equivariant endofunctor of V^G. Then the group G acts on $\mathbb{T}r(UFA')$ via the equivariance isomorphisms of id_V and UFA' (see Definition 5.2). In the situation where $F = \widetilde{H}$ for an equivariant endofunctor H of V, this action sends $\xi \in \mathbb{T}r(Hg)$ to



The action of G on $\mathbb{T}r(HA)$ in string diagram notation

In particular, the action on $\mathbb{T}r(A)$ agrees with that of the isomorphisms ψ_s, defined on page 25.

Consider the category $\mathrm{Vect_G}(G)$ of G-equivariant vectorbundles on G with respect to the conjugation action. The convolution product

$$(V \circledast W)_h := \bigoplus_{gs=h} V_g \otimes W_s$$

makes $\mathrm{Vect_G}(G)$ a braided monoidal category, where the braiding comes from the isomorphisms

$$V_g \otimes W_s \cong W_s \otimes V_{s^{-1}gs}$$

induced by ψ_s. A straight-forward manipulation of string diagrams[9] shows that R_ϱ is a commutative algebra object of $\mathrm{Vect}_{\mathrm{G}}(G)$ with respect to this (non-symmetric) braided monoidal structure.

Lemma 5.13. *The G-invariant part of the twisted group algebra R_ϱ is contained in the center of R_ϱ. More generally, let F be a 1-endomorphism of V^G, viewed as an equivariant 1-endomorphsim via* (11)*. Then we have*

$$\mathbb{T}r(M)^G \subseteq \mathrm{Center}_{\mathbb{T}r(A)}(\mathbb{T}r(M)).$$

Proof : Using the string diagram moves [Bar08, (2), (3), Lemma 6(i)], one sees that μ is in $\mathbb{T}r(M)^G$ if and only if for each $s \in G$ the following two string diagrams are equal:





It follows that μ is in the center of $\mathbb{T}r(M)$. □

Lemma 5.14. *Let F be an equivariant endofunctor of V^G. Then the isomorphism $j_{\widetilde{H}}$ of Lemma 5.11 induces an isomorphism of the G-invariant parts*

$$\mathbb{T}r(UFA')^G = 2\,\mathrm{Hom}_G(\mathrm{id}_V, UFA') \cong 2\,\mathrm{Hom}_{\mathrm{G}}(U, UF).$$

Corollary 5.15. *Let H be an equivariant 1-endomorphism of V. Then we have an isomorphism*

$$2\,\mathrm{End}_{\mathrm{G}}(U, HU) \cong \left(\bigoplus_{g\in G} \mathbb{T}r(Hg)\right)^G,$$

where the G-action on the right-hand side is as on Page 35.

---

[9] See [Bar08, (2), (3), Lemma 6(i)] for the relevant moves.

5.4 Traces in V^G

Throughout this section, we assume that we are given a linear 2-representation of G on V such that (V^G, U) exist.

Definition 5.16. Let W be an object of a 2-category $\mathcal{C}$. Then the *dimension* or *center* of W is the categorical trace of the identity 1-morphism of W,

$$\mathcal{D}\mathrm{im}(W) := \mathbb{T}r(\mathrm{id}_W) = 2\,\mathrm{Hom}(\mathrm{id}_W, \mathrm{id}_W).$$

Horizontal and vertical composition agree on $\mathcal{D}\mathrm{im}(W)$ and make it into a commutative monoid. If $\mathcal{C}$ is k-linear then $\mathcal{D}\mathrm{im}(W)$ is a commutative k-algebra.

Theorem 5.17. *We have an isomorphism of k-algebras*

$$i\colon \mathcal{D}\mathrm{im}(V^G) \xrightarrow{\cong} \left(\bigoplus_{g\in G} \mathbb{T}r(g)\right)^G.$$

For any 1-endomorphism F of V^G we have a bimodule isomorphism

$$i_F\colon \mathbb{T}r(F) \xrightarrow{\cong} \mathbb{T}r(UFA')^G.$$

Proof : We have

$$\begin{aligned}
\mathbb{T}r(F) &= 2\,\mathrm{Hom}\left(\mathrm{id}_{V^G}, F\right)\\
&= 2\,\mathrm{Hom}_{\mathrm{G}}\left(\mathrm{id}_{V^G}, F\right)\\
&\cong 2\,\mathrm{Hom}_{\mathrm{G}}^{\mathrm{triv}}\left(UF, UF\right)\\
&= 2\,\mathrm{Hom}_{\mathrm{G}}\left(UF, UF\right)\\
&\cong \mathbb{T}r(UFA')^G,
\end{aligned}$$

where the second equality is (11), the third isomorphism is the universal property of (V^G, U), the next equality is (12), and the last equality is Lemma 5.14. If $F = \mathrm{id}_{V^G}$ we may apply Corollary 5.15 to identify the last term with

$$\left(\bigoplus_{g\in G} \mathbb{T}r(g)\right)^G.$$

Explicitly, the isomorphism i in the statement of the theorem is given by

$$i\colon \zeta \longmapsto (U\zeta A') \circ \iota.$$

The fact that i preserves the algebra multiplication is proved by the equality of the following two string diagrams:



The multiplication of $i(\zeta)$ and $i(\vartheta)$ in $\mathbb{T}r(A)$ equals $i(\zeta\vartheta)$.

A similar identity of string diagrams shows that i_F is a map of bimodules. ☐

Corollary 5.18. *Let ϱ be a representation on an oject V of a linear 2-category, and assume that V^G exists. Then we have*

$$\dim\big(\mathcal{D}\mathrm{im}(V^G)\big) = \frac{1}{|G|}\sum_{gh=hg}\chi_\varrho(g,h).$$

Proof : We have

$$\begin{aligned}\dim\big(\mathbb{T}r(A)^G\big) &= \dim\bigoplus_{[g]\subseteq G}\mathbb{T}r(g)^{C_g}\\ &= \sum_{[g]}\frac{1}{|C_g|}\mathrm{tr}(h\circlearrowleft\mathbb{T}r(g))\\ &= \frac{1}{|G|}\sum_{gh=hg}\chi_\varrho(g,h).\end{aligned}$$

☐

Lemma 5.19. *Let $(H,\{\tau_g\})$ be an equivariant 1-endomorphism of V, and let $(\widetilde{H},\{\widetilde{\tau}_g\})$ be such that $U\widetilde{H}$ and HU agree as trivialized equivariant 1-morphisms. Then we have*

$$\widetilde{\tau}_g \cong \epsilon_g\widetilde{H}\epsilon_g^{-1}$$

(horizontal composition, vertical inverse). In other words, $\{\widetilde{\tau}_g\}$ agrees with the trivialization (11).

Corollary 5.20. *Let* $(H,\{\eta_g\})$ *is an equiariant 1-endomorphism of* V *and let* $(\widetilde{H},\{\widetilde{\eta}_g\})$ *then*

$$\mathbb{T}r(\widetilde{H}) \cong \left(\bigoplus_{g\in G} \mathbb{T}r(Hg)\right)^G.$$

5.5 Inner products

Definition 5.21. Let ϱ and π be 2-representations of G in a lax monoidal linear 2-category $(\mathcal{C},\boxtimes,1)$,and assume that the object $(\varrho\boxtimes\pi)^G$ exists. Then we define the inner product of ϱ and π to be the k-vector space

$$\langle\varrho,\pi\rangle_G := \mathcal{D}\mathrm{im}\left((\varrho\boxtimes\pi)^G\right).$$

As an immediate consequence of the corollary, we obtain

$$\dim_k\langle\varrho,\pi\rangle_G = \frac{1}{|G|}\sum_{gh=hg}\chi_\varrho(g,h)\cdot\chi_\pi(g,h).$$

6 Applications

6.1 Projective representations

Let $\mathcal{V} = \mathrm{Vect}_{\mathbb{C}}$. Then a linear G-action on $\mathcal{V}$ is classified by a 2-cocycle

$$c\colon G\times G\to\mathbb{C},$$

and $\mathcal{V}^G$ is identified with the category of projective representations of G^{op} with central charge

$$c^{op}(g,h) := c(h,g).$$

These are pairs

$$(W,\varphi\colon G^{op}\to\mathrm{Aut}(W)),$$

where W is a k-vector space, and φ is a map satisfying

$$\varphi(gh) = c(g,h)\cdot\varphi(h)\circ\varphi(g).$$

In particular, $\mathcal{V}^G$ is a 2-vectorspace, and

$$\dim\mathcal{D}\mathrm{im}(\mathcal{V}^G)$$

is the number of isomorphism classes of irreducible projective representations of G^{op} with central charge c^{op}.

On the other hand,

$$\mathbb{T}r(A)^G \cong \bigoplus_{[g]} \mathbb{C}^{C_g},$$

where the sum is over the conjugacy classes of G. The action of $h \in C_g$ on $\mathbb{C} \cong \mathbb{T}r(g)$ is multiplication with $\chi_{\varrho_c}(g,h)$.

Lemma 6.1. *We have*

$$\chi_{\varrho_c}(g,h) = \frac{c(h,g)}{c(g,h)}.$$

Proof : Apply the cocycle condition to $\delta(g,h,h^{-1})$ to get

$$c(gh,h^{-1})c(h,h^{-1})^{-1} = c(g,h)^{-1}c(g,1).$$

Substituting this into the formula of [GK08, Prop.5.1]), we obtain

$$\begin{aligned}\chi_{\varrho_c}(g,h) &= c(h,g)c(g,h)^{-1}c(g,1)c(1,1)^{-1}\\ &= c(h,g)c(g,h)^{-1}.\end{aligned}$$

□

A conjugacy class $[g] \subseteq G$ is called *c-regular* if

$$(\forall h \in C_g) \quad (c(g,h) = c(h,g)).$$

Note $[g]_G$ is c-regular if and only if $[g]_{G^{op}}$ is c^{op}-regular. Hence Theorem 5.17 specializes to the following result of Schur's:

Theorem 6.2 (Schur)**.** *The number of isomorphisms classes of irreducible projective representations with multiplier c equals the number of c-regular conjugacy classes of G.*

6.2 Algebras

Let A be an associative and unitary, finite dimensional k-algebra. Assume that G acts on A from the left by (unit preserving) algebra automorphisms. Let $\mathcal{V}$ = A-mod$^{\mathrm{f}}$ be as in Section 2.2.2. Then G acts on $\mathcal{V}$ from the right via the right-exact functors

$$M \longmapsto Mg.$$

Here Mg is the right A-module that is isomorphic to M as a k-vector space and whose A-action is twisted by g as follows:

$$(mg) \cdot a = (m \cdot g(a))g.$$

Write $A \rtimes G$ for the crossed product algebra. So,

$$A \rtimes G = \bigoplus_{g \in G} Ag$$

with multiplication

$$(a_1 g_1) \cdot (a_2 g_2) = (a_1 \cdot g_1(a_2))(g_1 g_2).$$

The category of (right-)equivariant objects in $\mathcal{V}$ is

$$\mathcal{V}^G \simeq (\mathrm{A} \rtimes \mathrm{G})\text{-}\mathrm{mod}^{\mathrm{f}}.$$

The twisted group algebra R_ϱ equals

$$\bigoplus_{g \in G} \mathrm{Center}_A(Ag).$$

Hence Theorem 5.17 becomes the well known formula

$$\mathrm{Center}(A \rtimes G) \cong \bigoplus_{[g]} \left(\mathrm{Center}_A(Ag)\right)^{C_g}. \tag{14}$$

6.3 Bimodules

Let A be an associative and unitary k-algebra, acted upon by G as in the previous section. View A as object of the 2-category $\mathcal{B}\mathrm{im}_k$ of Section 3.1. Let $U = A \rtimes G$, viewed as ($G \times G$-equivariant) $A \rtimes G$-A-bimodule. Under the equivalence (5) U corresponds to the forgetful functor sending an equivariant A-module to the underlying A-module. We would like to argue that the pair $(A \rtimes G, U)$ satisfies the universal property of Definition 5.6. This is not entirely true, but holds for a sufficiently large class of test-objects: let B be a second associative and unitary k-algebra on which G acts by algebra automorphisms.[10] Then composition with U gives an equivalence of categories

$$\mathcal{B}\mathrm{im}_{k,G \times G}(B, A) \simeq \mathcal{B}\mathrm{im}_{k,G}(B, A \rtimes G).$$

---

[10] For the full universal property to by satisfied, we would need to allow G to act by 1-automorphisms.

This is sufficient to ensure that the proof of Theorem 5.17 goes through, again yielding (14).

We now view A and $A \rtimes G$ as objects of the 2-category $\mathcal{DB}\mathrm{im}_{\mathrm{k}}$ (see Section 3.2), and we write $\mathcal{U}$ for the complex consisting of U situated in degree zero and zeros elsewhere. For a test-object B as above, we need to distinguish between the derived category of equivariant bimodules and the category of equivariant 1-morphisms: in general, the forgetful functor

$$\mathcal{DB}\mathrm{im}_{k,G}(B, A) \longrightarrow \mathcal{DB}\mathrm{im}(B, A)^G = 1\text{-}\mathrm{Hom}_{\mathrm{G}}(B, A),$$

is not an equivalence of categories.

In addition to restricting our pool of test-objects as above, we need to modify the universal property in Definition 5.6, replacing $1\text{-}\mathrm{Hom}_G$ with $\mathcal{DB}\mathrm{im}_{\mathrm{k,G}}$. With this modification to Theorem 5.17 and its proof,[11] we obtain the isomorphism

$$\mathrm{HH}^\bullet(A \rtimes G, A \rtimes G) \cong \left(\bigoplus_{g \in G} \mathrm{HH}^\bullet(A, Ag) \right)^G$$

of [DE05, Prop 3].[12]

6.4 Coherent sheaves

Let X be a smooth projective variety over k, let $\mathcal{C}\mathrm{oh}(X)$ be the category of coherent sheaves on X, and assume that a finite group G acts on X from the left. Then $g \in G$ acts on $\mathcal{C}\mathrm{oh}(X)$ via

$$\mathcal{F} \longmapsto g_* \mathcal{F}, \tag{15}$$

and the category of equivariant objects in $\mathcal{C}\mathrm{oh}(X)$ is identified with the category of G-equivariant sheaves on X,

$$\mathcal{C}\mathrm{oh}(X)^G \simeq \mathcal{C}\mathrm{oh}_{\mathrm{G}}(X).$$

We may view $\mathcal{C}\mathrm{oh}_{\mathrm{G}}(X)$ as the category of sheaves on the orbifold quotient $[X/G]$.

---

[11] The last step of the proof is Lemma A.7.

[12] It appears that this result goes back to an unpublished preprint by Brylinski, dating from 1987.

It is possible to extend the 2-category of kernels $\mathcal{V}\mathrm{ar}$ of Section 3.4 so that it contains both X and $[X/G]$ as objects (see [CW10, p.6] or [Căl]). The formalism remains the same as in Section 3.4. Recall that the G-action (15) on $\mathcal{D}^+\mathcal{C}\mathrm{oh}(X)$ lifts to a 2-representation

$$\varrho\colon G \longrightarrow 1\mathrm{Hom}(X, X)$$

in $\mathcal{V}\mathrm{ar}$, given by the kernels $\varrho(g) = \mathscr{O}_{\Gamma_g}$.

Let Y be a second proper, smooth space with G-action, and write $\sigma(s) := \mathscr{O}_{\Gamma_s}$ for the corresponding 2-representation of kernels on Y. Using the projection formula and flat base-change (c.f. [Căl05, 2.7]), one finds natural isomorphisms

$$\varrho(g) \circ \mathcal{K} \cong R(1, g)_* \mathcal{K}$$

and

$$\mathcal{K} \circ \sigma(s^{-1}) \cong L(s^{-1}, 1)^* \mathcal{K} \cong R(s, 1)_* \mathcal{K}.$$

So,

$$\varrho(g) \circ \mathcal{K} \circ \sigma(s^{-1}) \cong R(s, g)_* \mathcal{K},$$

and the categories of equivariant 1-homomorphisms of Definition 5.2 are identified as

$$1\mathrm{Hom}_{\mathrm{G}}(Y, X) \simeq \left(\mathcal{D}^+\mathcal{C}\mathrm{oh}(Y \times X)\right)^G$$

and

$$1\mathrm{Hom}^{\mathrm{triv}}_{\mathrm{G}}(Y, X) \simeq \left(\mathcal{D}^+\mathcal{C}\mathrm{oh}(Y \times X)\right)^{G\times G}.$$

As in the previous section, we need to replace these with

$$\mathcal{D}^+\mathcal{C}\mathrm{oh}_{\mathrm{G}}(Y \times X) \quad \text{and} \quad \mathcal{D}^+\mathcal{C}\mathrm{oh}_{\mathrm{G}\times\mathrm{G}}(Y \times X).$$

With this modification, $[X/G]$ satisfies the universal property for test-objects Y as above,

$$\mathcal{D}^+\mathcal{C}\mathrm{oh}_{\mathrm{G}}(Y \times [X/G]) \simeq \mathcal{D}^+\mathcal{C}\mathrm{oh}_{\mathrm{G}\times\mathrm{G}}(Y \times X),$$

and the proof of Theorem 5.17 goes through (using Lemma A.7) to yield

$$\mathrm{HH}^\bullet([X/G]) \cong \bigoplus_{[g]} \mathbb{T}r^\bullet(g)^{C_g}.$$

The right-hand side of this isomorphism is identified by Theorem 3.1. We have proved

Theorem 6.3. *We have*

$$\mathrm{HH}^\bullet([X/G]) \cong \left(\bigoplus_{g,\alpha} \mathrm{HH}^{\bullet - \mathrm{codim}(\mathrm{X}^{\mathrm{g}}_{\alpha})} \left(X^g_\alpha, \det(N^g)\right) \right)^G ,$$

where α runs over the connected components of the fixed point sets.

6.5 The example of a trivial G-action

Let $\mathcal{B}$ be a k-linear abelian category, and assume that G acts trivially (from the left) on $\mathcal{B}$. Then the data of an equivariant object

$$(B, \epsilon) \in \mathrm{ob}\mathcal{B}^G$$

are equivalent to that of an object $B \in \mathrm{ob}\mathcal{B}$ together with a right G-action

$$\epsilon\colon k[G] \to \mathrm{End}(B).$$

By [Del90, 5.11], we have an equivalence of categories

$$\begin{aligned} \mathcal{B}^G &\simeq \left(\mathrm{k[G]\text{-}mod^f}\right) \widehat{\boxtimes} \mathcal{B} \\ &\simeq (W_1 \boxtimes \mathcal{B}) \boxplus \cdots \boxplus (W_r \boxtimes \mathcal{B}) \\ &\simeq \left(\mathrm{k[G]\text{-}mod^f}\right) \boxtimes \mathcal{B}, \end{aligned}$$

where the W_i represent the isomorphism classes of irreducible representations of G. Here third equivalence follows from the construction of $\boxtimes$, and the second equivalence holds, because the category in the second row is already abelian.

Example 6.4. Let $\mathcal{B}$ be $\mathcal{B} = \mathrm{Vect}^{\mathbb{C}}(X)$. Then $\mathcal{B}^G = \mathrm{Vect}^{\mathbb{C}}_{\mathrm{G}}(X)$ is the category of vector bundles on X with fibre-preserving G-action. In this case the above equivalences boil down to the well known fact that any G-vector bundle V over a base-space with trivial G-action can be decomposed as

$$V \cong \bigoplus_i W_i \otimes V_i,$$

where the $V_i \in \mathrm{Vect}^{\mathbb{C}}(X)$ are non-equivariant vector bundles.

We have seen two ways to calculate the center of $\mathcal{B}^G$. By Theorem 2.5,

$$\mathcal{D}\mathrm{im}\left(\mathcal{B}^G\right) \cong \mathrm{Center}(k[G]) \otimes \mathcal{D}\mathrm{im}(\mathcal{B}).$$

On the other hand, since G acts trivially on $\mathcal{B}$, Theorem 5.17 becomes

$$\begin{aligned} \mathcal{D}\mathrm{im}\left(\mathcal{B}^G\right) &\cong \bigoplus_{[g]} \mathbb{T}r(g)^{C_g} \\ &\cong \bigoplus_{[g]} \mathcal{D}\mathrm{im}(\mathcal{B}). \end{aligned}$$

Indeed, these two results agree. In the special case where $\mathcal{B} = \mathrm{Vect}_{\mathrm{k}}$, we can be more specific and identify the isomorphism

$$i\colon \mathrm{Center}(k[G]) \longrightarrow \bigoplus_{[g]} k$$

of Theorem 5.17: its inverse i^{-1} sends the $[g]$th basis vector to the element

$$e_{[g]} \coloneqq \sum_{h \in C_g} h^{-1}$$

of $\mathrm{Center}(k[G])$.

A Clean intersections

This appendix collects some results about clean intersections that are probably well-known to the experts. I did not know references, so I am including proofs.

Lemma A.1. *Let $W \subseteq Z$ be a closed subscheme defined by the ideal sheaf $\mathcal{I} \subseteq \mathscr{O}_Z$. Let $w \in W$, and write $\mathfrak{m}$ for the maximal ideal of $\mathscr{O}_{Z,w}$. Then the conormal space of W in Z at w is naturally identified with the quotient*

$$N_w^\vee \cong \mathcal{I}_w / \left(\mathcal{I}_w \cap \mathfrak{m}^2\right).$$

If $\mathcal{I}_w$ is a prime ideal in $\mathscr{O}_{Z,w}$, this simplifies to

$$N_w^\vee \cong \mathcal{I}_w / \mathfrak{m}\mathcal{I}_w.$$

Proof : Write $\mathfrak{n}$ for the maximal ideal of

$$\mathscr{O}_{W,w} = \mathscr{O}_{Z,w}/\mathcal{I}_w.$$

Then $\mathfrak{n} = \mathfrak{m}/\mathcal{I}_w$, and we have the short exact sequences

$$\begin{tikzcd}
0 \arrow[r] & \mathcal{I}_w/\mathcal{I}_w \cap \mathfrak{m}^2 \arrow[r] \arrow[d] & \mathfrak{m}/\mathfrak{m}^2 \arrow[r] \arrow[d] & \mathfrak{n}/\mathfrak{n}^2 \arrow[r] \arrow[d] & 0 \\
0 \arrow[r] & N_w^\vee \arrow[r] & T_{Z,w}^\vee \arrow[r] & T_{W,w}^\vee \arrow[r] & 0.
\end{tikzcd}$$

Hence the first claim. If $\mathcal{I}_w$ is prime, then

$$\mathcal{I}_w \cap \mathfrak{m}^2 = \mathfrak{m}\mathcal{I}_w,$$

hence the second claim. ☐

Corollary A.2. *Let $i\colon W \hookrightarrow Z$ be the inclusion of an irreducible closed subscheme, and let $w \in W$. If*

$$T_{i,w}\colon T_{W,w} \longrightarrow T_{Z,w}$$

is an isomorphism then i is an insomorphism in a neighbourhood of w.

Proof : The map $T_{i,w}$ is an isomorphism if and only if $N_w^\vee = \{0\}$. Since W is irreducible, this is equivalent to $I_w = \mathfrak{m}\mathcal{I}_w$. By Nakayama's Lemma, this is equivalent to $\mathcal{I}_w = 0$. ☐

Lemma A.3. *Let $i\colon W \hookrightarrow Z$ be the inclusion of a closed subscheme, and let $w \in W$. Then there exists an open neighbourhood U of w in Z and a regular closed subscheme $Y \subseteq U$ such that Y contains $W \cap U$ and $T_{Y,w} = T_{W,w}$ inside $T_{Z,w}$.*

Proof : Let $\mathfrak{m}$, $\mathfrak{n}$, and $\mathcal{I}$ be as in Lemma A.1. Pick a basis $f_1, \ldots, f_d$ of

$$N_w^\vee \cong \mathcal{I}_w/\left(\mathcal{I}_w \cap \mathfrak{m}^2\right).$$

By [Liu02, Cor 4.2.12], this sequence is regular. Hence we can find an open neighbourhood U of w in Z and a regular sequence

$$\widetilde{f}_1, \ldots, \widetilde{f}_d \in \Gamma(U, \mathcal{I}),$$

representing $f_1, \dots, f_d$. By construction, we have $W \subseteq Y$, the closed immersion $Y \hookrightarrow U$ is regular at w, and the conormal spaces of W and Y inside Z at w agree. $\square$

Lemma A.4 ([Li09, Lem.5.1])**.** *Let Z be a nonsingular variety, and let X and Y be nonsingular closed subvarieties of Z. Then the connected components of the scheme-theoretic intersection $X \cap Y$ of X and Y in Z are non-singular if and only if the following differential geometric conditions are satisfied:*

1. *the connected components of the "set-theoretic intersection"*

$$(X \cap Y)_{red}$$

   are nonsingular varieties, and

2. *we have*

$$T_X \cap T_Y = T_{X \cap Y}$$

   inside T_Z.

In the context of differential geometry, these conditions define the notion of *clean intersection*, due to Bott [Bot56].

Lemma A.5. *In the situation of the previous Lemma the canonical map of vector bundles over $X \cap Y$*

$$T_X + T_Y \hookrightarrow T_{X \cup Y}$$

is an isomorphism. Here the sum on the left-hand side is taken inside $T_Z|_{X \cap Y}$.

Proof : Fix $z \in X \cap Y$, let $\mathfrak{m} \subset \mathscr{O}_{Z,z}$ be the maximal ideal, and let $\mathcal{I}$ and $\mathcal{J}$ denote the stalks of the ideal sheaves of X and Y at z. Then we have a cartesian square

$$\begin{tikzcd}
\mathcal{I} \cap \mathcal{J} / \mathcal{I} \cap \mathcal{J} \cap \mathfrak{m}^2 \arrow[r, tail] \arrow[d, tail] & \mathcal{I} / \mathcal{I} \cap \mathfrak{m}^2 \arrow[d, tail] \\
\mathcal{J} / \mathcal{J} \cap \mathfrak{m}^2 \arrow[r, tail] & \mathcal{I} + \mathcal{J} / (\mathcal{I} + \mathcal{J}) \cap \mathfrak{m}^2.
\end{tikzcd}$$

Lemma A.1 identifies the vector spaces in this diagram with the conormal spaces of $X \cup Y$, X, Y, and $X \cap Y$ inside Z at z. A dimension count now proves the claim. $\square$

Lemma A.6. *Let Z be a nonsingular variety, and let X, Y, and W be nonsingular closed subvarieties of Z such that W is contained in X and in Y ("contained" means as a closed subscheme). Let $w \in W$, and assume that the map*

$$T_{W,w} \hookrightarrow T_{X,w} \cap T_{Y,w}$$

is an isomorphisms. (Here the intersection on the right-hand side is inside $T_{Z,z}$.) Then there exists a neighbourhood of w in Z inside which the map $i\colon W \hookrightarrow X \cap Y$ is an isomorphism.

Proof : We know that the composite

$$\begin{tikzcd} T_{W,w} \arrow[r, tail, "T_{i,w}"] & T_{X\cap Y} \arrow[r, tail] & T_X \cap T_Y \end{tikzcd}$$

is an isomorphism. Hence $T_{i,w}$ is an isomorphism, as well. Applying Corollary A.2, we obtain the claim. □

We will also need a lemma from homological algebra.

Lemma A.7. *Let $\mathcal{A}$ be an abelian category, and assume that $\mathcal{A}^G$ has enough injectives. Let $\mathcal{F}$ and $\mathcal{G}$ be equivariant objects in $\mathcal{A}$. Then we have*

$$\mathrm{Hom}^{\bullet}_{\mathcal{D}^+_G(\mathcal{A})}(\mathcal{F},\mathcal{G}) \cong \mathrm{Hom}^{\bullet}_{\mathcal{D}^+(\mathcal{A})}(\mathcal{F},\mathcal{G})^G.$$

Proof : Choose an injective resolution $\mathcal{G} \to \mathcal{I}^\bullet$ in $\mathcal{A}^G$. Since the forgetful functor $U\colon \mathcal{A}^G \to \mathcal{A}$ is a right-adjoint, it preserves injectives. Therefore, we have

$$\begin{aligned} \mathrm{Hom}^{\bullet}_{\mathcal{D}^+_G(\mathcal{A})}(\mathcal{F},\mathcal{G}) &\cong \mathrm{Hom}^{\bullet}_{\mathcal{C}h^+_G(\mathcal{A})}(\mathcal{F},\mathcal{I}^\bullet) \\ &\cong \mathrm{Hom}^{\bullet}_{\mathcal{C}h^+(\mathcal{A})}\left(\mathcal{F},\mathcal{I}^\bullet\right)^G \\ &\cong \mathrm{Hom}^{\bullet}_{\mathcal{D}^+(\mathcal{A})}\left(\mathcal{F},\mathcal{G}\right)^G. \end{aligned}$$

□

References

[Bar03] Vladimir Baranovsky. Orbifold cohomology as periodic cyclic homology. *Internat. J. Math.*, 14(8):791–812, 2003.

[Bar08] Bruce Bartlett. *On unitary 2-representations of finite groups and topological quantum field theory.* PhD thesis, University of Sheffield, 2008. arXiv:0901.3975v1.

[BK01] Bojko Bakalov and Alexander Kirillov, Jr. *Lectures on tensor categories and modular functors*, volume 21 of *University Lecture Series.* American Mathematical Society, Providence, RI, 2001.

[BL03] John C. Baez and Laurel Langford. Higher-dimensional algebra. IV. 2-tangles. *Adv. Math.*, 180(2):705–764, 2003.

[BLL04] Alexey I. Bondal, Michael Larsen, and Valery A. Lunts. Grothendieck ring of pretriangulated categories. *Int. Math. Res. Not.*, (29):1461–1495, 2004.

[BM06] John W. Barrett and Marco Mackaay. Categorical representations of categorical groups. *Theory Appl. Categ.*, 16:No. 20, 529–557 (electronic), 2006.

[BN96] John C. Baez and Martin Neuchl. Higher-dimensional algebra. I. Braided monoidal 2-categories. *Adv. Math.*, 121(2):196–244, 1996.

[Bot56] Raoul Bott. On the iteration of closed geodesics and the Sturm intersection theory. *Comm. Pure Appl. Math.*, 9:171–206, 1956.

[BWC07] John C. Baez, Derek K. Wise, and Alissa S. Crans. Exotic statistics for strings in 4D BF theory. *Adv. Theor. Math. Phys.*, 11(5):707–749, 2007.

[BZ07] David Ben-Zvi. Lecture series on the geometric Langlands correspondence. Mathematical Institute, Oxford, April 2007.

[Căl] Andrei Căldăraru. The Mukai pairing, iii, Calabi-Yau orbifolds.

[Căl05] Andrei Căldăraru. Derived categories of sheaves: a skimming. In *Snowbird lectures in algebraic geometry*, volume 388 of *Contemp. Math.*, pages 43–75. Amer. Math. Soc., Providence, RI, 2005.

[CKS03] Andrei Căldăraru, Sheldon Katz, and Eric Sharpe. D-branes, B fields, and Ext groups. *Adv. Theor. Math. Phys.*, 7(3):381–404, 2003.

[Cra98] Sjoerd E. Crans. Generalized centers of braided and sylleptic monoidal 2-categories. *Adv. Math.*, 136(2):183–223, 1998.

[CW10] Andrei Căldăraru and Simon Willerton. The Mukai pairing. I. A categorical approach. *New York J. Math.*, 16:61–98, 2010.

[CY05] Louis Crane and David N. Yetter. Measurable categories and 2-groups. *Appl. Categ. Structures*, 13(5-6):501–516, 2005.

[DE05] Vasiliy Dolgushev and Pavel Etingof. Hochschild cohomology of quantized symplectic orbifolds and the Chen-Ruan cohomology. *Int. Math. Res. Not.*, (27):1657–1688, 2005.

[Del90] P. Deligne. Catégories tannakiennes. In *The Grothendieck Festschrift, Vol. II*, volume 87 of *Progr. Math.*, pages 111–195. Birkhäuser Boston, Boston, MA, 1990.

[Del97] P. Deligne. Action du groupe des tresses sur une catégorie. *Invent. Math.*, 128(1):159–175, 1997.

[Elg07] Josep Elgueta. Representation theory of 2-groups on Kapranov and Voevodsky's 2-vector spaces. *Adv. Math.*, 213(1):53–92, 2007.

[Fre94] Daniel S. Freed. Higher algebraic structures and quantization. *Comm. Math. Phys.*, 159(2):343–398, 1994.

[Gan06] Nora Ganter. Orbifold genera, product formulas and power operations. *Adv. Math.*, 205(1):84–133, 2006.

[GK] Nora Ganter and Mikhail Kapranov. Symmetric and exterior powers of categories. in preparation.

[GK04] Victor Ginzburg and Dmitry Kaledin. Poisson deformations of symplectic quotient singularities. *Adv. Math.*, 186(1):1–57, 2004.

[GK08] Nora Ganter and Mikhail Kapranov. Representation and character theory in 2-categories. *Adv. Math.*, 217(5):2268–2300, 2008.

[Gro57] Alexander Grothendieck. Sur quelques points d'algèbre homologique. *Tôhoku Math. J. (2)*, 9:119–221, 1957.

[HKR00] Michael J. Hopkins, Nicholas J. Kuhn, and Douglas C. Ravenel. Generalized group characters and complex oriented cohomology theories. *J. Amer. Math. Soc.*, 13(3):553–594 (electronic), 2000.

[Huy06] D. Huybrechts. *Fourier-Mukai transforms in algebraic geometry.* Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, Oxford, 2006.

[Kho00] Mikhail Khovanov. A categorification of the Jones polynomial. *Duke Math. J.*, 101(3):359–426, 2000.

[KS94] Masaki Kashiwara and Pierre Schapira. *Sheaves on manifolds*, volume 292 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences].* Springer-Verlag, Berlin, 1994. With a chapter in French by Christian Houzel, Corrected reprint of the 1990 original.

[KV94] M. M. Kapranov and V. A. Voevodsky. 2-categories and Zamolodchikov tetrahedra equations. In *Algebraic groups and their generalizations: quantum and infinite-dimensional methods (University Park, PA, 1991)*, volume 56 of *Proc. Sympos. Pure Math.*, pages 177–259. Amer. Math. Soc., Providence, RI, 1994.

[Li09] Li Li. Wonderful compactification of an arrangement of subvarieties. *Michigan Math. J.*, 58(2):535–563, 2009.

[Lip09] Joseph Lipman. *Notes on derived functors and Grothendieck duality. Foundations of Grothendieck duality for diagrams of schemes.* Number 1960 in Lecture Notes in Mathematics. Springer Verlag, New York, 2009.

[Liu02] Qing Liu. *Algebraic geometry and arithmetic curves*, volume 6 of *Oxford Graduate Texts in Mathematics.* Oxford University Press, Oxford, 2002. Translated from the French by Reinie Erné, Oxford Science Publications.

[Lus87] George Lusztig. Introduction to character sheaves. In *The Arcata Conference on Representations of Finite Groups (Arcata, Calif., 1986)*, volume 47 of *Proc. Sympos. Pure Math.*, pages 165–179. Amer. Math. Soc., Providence, RI, 1987.

[ML95] Saunders Mac Lane. *Homology*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1975 edition.

[Oso10] Angélica M. Osorno. Explicit formulas for 2-characters. *Topology Appl.*, 157(2):369–377, 2010.

[Ost03] Victor Ostrik. Module categories, weak Hopf algebras and modular invariants. *Transform. Groups*, 8(2):177–206, 2003.

[Ser77] Jean-Pierre Serre. *Linear representations of finite groups*. Springer-Verlag, New York, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42.

[ST01] Paul Seidel and Richard Thomas. Braid group actions on derived categories of coherent sheaves. *Duke Math. J.*, 108(1):37–108, 2001.

[Str00] N. P. Strickland. $K(N)$-local duality for finite groups and groupoids. *Topology*, 39(4):733–772, 2000.